

Sinclairville Elementary Health Office

News and Alerts

May is Hand Hygiene Month

May 1 2023



How Germs Spread

Washing hands can keep you healthy and prevent the spread of respiratory and diarrheal infections. Germs can spread from person to person or from surfaces to people when you:

- Touch your eyes, nose, and mouth with unwashed hands
- Prepare or eat food and drinks with unwashed hands
- Touch surfaces or objects that have germs on them
- Blow your nose, cough, or sneeze into hands and then touch other people's hands or common objects

Key Times to Wash Hands

You can help yourself and your loved ones stay healthy by washing your hands often, especially during these key times when you are likely to get and spread germs:

- Before, during, and after preparing food
- Before and after eating food
- Before and after caring for someone at home who is sick with vomiting or diarrhea
- Before and after treating a cut or wound
- After using the toilet
- After changing diapers or cleaning up a child who has used the toilet
- After blowing your nose, coughing, or sneezing
- After touching an animal, animal feed, or animal waste
- After handling pet food or pet treats
- After touching garbage

Use Hand Sanitizer When You Can't Use Soap and Water

You can use an alcohol-based hand sanitizer that contains at least 60% alcohol if soap and water are not available.

Washing hands with soap and water is the best way to get rid of germs in most situations. If soap and water are not readily available, you can use an alcohol-based hand sanitizer that contains at least 60% alcohol. You can tell if the sanitizer contains at least 60% alcohol by looking at the product label.

Sanitizers can quickly reduce the number of germs on hands in many situations. However,

- Sanitizers do not get rid of all types of germs.
- Hand sanitizers may not be as effective when hands are visibly dirty or greasy.
- Hand sanitizers might not remove harmful chemicals from hands like pesticides and heavy metals.

Caution! Swallowing alcohol-based hand sanitizers can cause alcohol poisoning if more than a couple of mouthfuls are swallowed. Keep it out of reach of young children and supervise their use.

How to Use Hand Sanitizer

1. Apply the gel product to the palm of one hand (read the label to learn the correct amount).
2. Rub your hands together.
3. Rub the gel over all the surfaces of your hands and fingers until your hands are dry. This should take around 20 seconds.

More Information

- <https://www.cdc.gov/handwashing/faqs.html>
- <https://www.cdc.gov/handhygiene/index.html>
- <https://www.cdc.gov/handwashing/training-education.html>
- <https://www.cdc.gov/handwashing/global-handwashing-day.html>
- <http://www.nimh.nih.gov/health/topics/attention-deficit-hyperactivity-disorder-adhd/index.shtml>

Brought to you by:

Samantha Smith RT (R), RN

Elementary School Nurse

Jacqueleen Anderson LPN

Elementary School Nurse

