
2

The Chemical Context of Life

Key Concepts

- 2.1 Matter consists of chemical elements in pure form and in combinations called compounds
- 2.2 An element's properties depend on the structure of its atoms
- 2.3 The formation and function of molecules depend on chemical bonding between atoms
- 2.4 Chemical reactions make and break chemical bonds
- 2.5 Hydrogen bonding gives water properties that help make life possible on Earth



Overview

The Importance of Chemistry to Life

As the Earth is warming from climate change, its life support systems are changing. In the Arctic, sea ice and glaciers are melting, affecting life all around them. The warmer Arctic waters and smaller ice pack are resulting in blooms of phytoplankton

(microscopic aquatic photosynthetic organisms), whose various shades

of blue and green can be seen from space (Figure 2.1).

While phytoplankton are thriving, however, organisms that depend on Arcticice are suffering, such as the black guillemots shown here (also, see Figure 2.21).

■ Black guillemots, threatened by climate change Three-quarters of Earth's surface is covered by water, a remarkable substance with rare properties emerging from the chemistry of the water molecule. Water is the only common substance on Earth to exist in nature in all three physical states of matter: solid (ice), liquid water, and a gas (water vapor). Furthermore, the solid form of water floats on the liquid form, a rare, life-supporting property emerging from the chemistry of the water molecule.

Understanding the chemical characteristics of water and other substances is central to biology, the study of life. Biology is interdisciplinary: Organisms and their environments are natural systems to which the concepts of chemistry and physics apply. This unit of chapters introduces some basic concepts of chemistry that apply to the study of life. Somewhere in the transition from molecules to cells, we will cross the blurry boundary between nonlife and life. This chapter focuses on the chemical

components that make up all matter, with a final section on the substance that supports all of life—water.

Go to the Mastering Biology eText or Study Area.

Mastering Biology Get Ready for This Chapter

CONCEPT 2.1

Matter consists of chemical elements in pure form and in combinations called compounds

Organisms are composed of **matter**, which is anything that takes up space and has mass. Matter exists in many forms. Rocks, metals, oils, gases, and living organisms are a few examples of what seems to be an endless assortment of matter.

Elements and Compounds

Matter is made up of elements. An **element** is a substance that cannot be broken down to other substances by chemical reactions. Today, chemists recognize 92 elements occurring in nature; gold, copper, carbon, and oxygen are examples. Each element has a symbol, usually the first letter or two of its name. Some symbols are derived from Latin or German; for instance, the symbol for sodium is Na, from the Latin word *natrium*.

A **compound** is a substance consisting of two or more different elements combined in a fixed ratio. Table salt, for example, is sodium chloride (NaCl), a compound composed of the elements sodium (Na) and chlorine (Cl) in a 1:1 ratio. Pure sodium is a metal, and pure chlorine is a poisonous gas. When chemically combined, however, sodium and chlorine form an edible compound. Water (H_2O), another compound, consists of the elements hydrogen (H) and oxygen (O) in a 2:1 ratio. These are simple examples of organized matter having *emergent properties*: A compound has chemical and physical characteristics different from those of its constituent elements (**Figure 2.2**).

The Elements of Life

Of the 92 natural elements, about 20–25% are **essential elements** that an organism needs to live a healthy life and reproduce. The essential elements are similar among organisms, but there is some variation—for example, humans need 25 elements, but plants need only 17.

▼ Figure 2.2 The emergent properties of a compound. The metal sodium combines with the poisonous gas chlorine, forming the edible compound sodium chloride, or table salt.




Table 2.1 Elements in the Human Body

Element	Symbol	Percentage of Body Mass (including water)	
Oxygen	0	65.0%	
Carbon	С	18.5%	96.3%
Hydrogen	Н	9.5%	
Nitrogen	N	3.3%	
Calcium	Ca	1.5%	
Phosphorus	P	1.0%	
Potassium	K	0.4%	
Sulfur	S	0.3%	3.7%
Sodium	Na	0.2%	
Chlorine	Cl	0.2%	
Magnesium	Mg	0.1%	

Trace elements (less than 0.01% of mass): boron (B), chromium (Cr), cobalt (Co), copper (Cu), fluorine (F), iodine (I), iron (Fe), manganese (Mn), molybdenum (Mo), selenium (Se), silicon (Si), tin (Sn), vanadium (V), zinc (Zn)

INTERPRET THE DATA Given the makeup of the human body, what compound do you think accounts for the high percentage of oxygen?

Relative amounts of all the elements in the human body are listed in **Table 2.1**. Just four elements—oxygen (O), carbon (C), hydrogen (H), and nitrogen (N)—make up approximately 96% of living matter. Calcium (Ca), phosphorus (P), potassium (K), sulfur (S), and a few other elements account for most of the remaining 4% or so of an organism's mass. **Trace elements** are required by an organism in only minute quantities. Some trace elements, such as iron (Fe), are needed by all forms of life; others are required only by certain species. For example, in vertebrates (animals with backbones), the element iodine (I) is an essential ingredient of a hormone produced by the thyroid gland. A daily intake of only 0.15 milligram (mg) of iodine is adequate for normal activity of the human thyroid. An iodine deficiency in the diet causes the thyroid gland to grow to abnormal size, a condition called goiter. Consuming seafood or iodized salt reduces the incidence of goiter.

Evolution of Tolerance to Toxic Elements

EVOLUTION Some naturally occurring elements are toxic to organisms. In humans, for instance, the element arsenic has been linked to numerous diseases and can be lethal. Some species, however, have become adapted to environments containing elements that are usually toxic. For example, sunflower plants can take up lead, zinc, and other heavy metals in concentrations that would kill most organisms. This allowed scientists to use sunflowers to detoxify contaminated soils after Hurricane Katrina. Presumably, variants of ancestral sunflower species were able to grow in soils with heavy metals, and subsequent natural selection resulted in their survival and reproduction.

CONCEPT CHECK 2.1

- 1. Is a trace element an essential element? Explain.
- 2. WHAT IF? In humans, iron is a trace element required for the proper functioning of hemoglobin, the molecule that carries oxygen in red blood cells. What might be the effects of an iron deficiency?

For suggested answers, see Appendix A.

CONCEPT 2.2

An element's properties depend on the structure of its atoms

Each element consists of a certain type of atom that is different from the atoms of any other element. An **atom** is the smallest unit of matter that still retains the properties of an element. Atoms are so small that it would take about a million of them to stretch across the period at the end of this sentence. We symbolize atoms with the same abbreviation used for the element that is made up of those atoms. For example, the symbol C stands for both the element carbon and a single carbon atom.


Subatomic Particles

Although the atom is the smallest unit having the properties of an element, these tiny bits of matter are composed of even smaller parts, called *subatomic particles*. Using high-energy collisions, physicists have produced more than 100 types of particles from the atom, but only three kinds of particles are relevant here: **neutrons**, **protons**, and **electrons**. Protons and electrons are electrically charged. Each proton has one unit of positive charge, and each electron has one unit of negative charge. A neutron, as its name implies, is electrically neutral.

Protons and neutrons are packed together tightly in a dense core, or **atomic nucleus**, at the center of an atom; protons give the nucleus a positive charge. The rapidly moving electrons form a "cloud" of negative charge around the nucleus, and it is the attraction between opposite charges that keeps the electrons in the vicinity of the nucleus. **Figure 2.3** shows two commonly used models of the structure of the helium atom as an example.

The neutron and proton are almost identical in mass, each about $1.7 \times 10^{-24} \, \mathrm{gram}$ (g). Grams and other conventional units are not very useful for describing the mass of objects that are so minuscule. Thus, for atoms and subatomic particles (and for molecules, too), we use a unit of measurement called the **dalton** (the same as the *atomic mass unit*, or *amu*). Neutrons and protons have masses close to 1 dalton. Because the mass of an electron is only about 1/2,000 that of a neutron or proton, we can ignore electrons when computing the total mass of an atom.

▼ Figure 2.3 Simplified models of a helium (He) atom. The helium nucleus consists of 2 neutrons (brown) and 2 protons (pink). Two electrons (yellow) exist outside the nucleus. These models are not to scale; they greatly overestimate the size of the nucleus in relation to the electron cloud.



- (a) This model represents the two electrons as a cloud of negative charge, a result of their motion around the nucleus.
- (b) In this more simplified model, the electrons are shown as two small yellow spheres on a circle around the nucleus.

Atomic Number and Atomic Mass

Atoms of the various elements differ in their number of subatomic particles. All atoms of a particular element have the same number of protons in their nuclei. This number of protons, which is unique to that element, is called the **atomic number** and is written as a subscript to the left of the symbol for the element. The abbreviation ₂He, for example, tells us that an atom of the element helium has 2 protons in its nucleus. Unless otherwise indicated, an atom is neutral in electrical charge, which means that its protons must be balanced by an equal number of electrons. Therefore, the atomic number tells us the number of protons and also the number of electrons in an electrically neutral atom.

We can deduce the number of neutrons from a second quantity, the **mass number**, which is the total number of protons and neutrons in the nucleus of an atom. The mass number is written as a superscript to the left of an element's symbol. For example, we can use this shorthand to write an atom of helium as ${}_{2}^{4}$ He. Because the atomic number indicates how many protons there are, we can determine the number of neutrons by subtracting the atomic number from the mass number: In our example, the helium atom ${}_{2}^{4}$ He has 2 neutrons. For sodium (Na):

Number of neutrons = mass number – atomic number = 23 - 11 = 12 for sodium

The simplest atom is hydrogen ¹H, which has no neutrons; it consists of a single proton with a single electron.

Because the contribution of electrons to mass is negligible, almost all of an atom's mass is concentrated in its nucleus. Neutrons and protons each have a mass very close to 1 dalton, so the mass number is close to, but slightly different from, the total mass of an atom, called its **atomic mass**. For example, the mass number of sodium $\binom{23}{11}$ Na) is 23, but its atomic mass is 22.9898 daltons; the difference is explained below.

Mastering Biology
 Animation: Atomic Number and Atomic Mass

Isotopes

All atoms of a given element have the same number of protons, but some atoms have more neutrons than other atoms of the same element and thus have greater mass. These different atomic forms of the same element are called isotopes of the element. In nature, an element may occur as a mixture of its isotopes. As an example, the element carbon, which has the atomic number 6, has three naturally occurring isotopes. The most common isotope is carbon-12, ¹²₆C, which accounts for about 99% of the carbon in nature. The isotope ¹²C has 6 neutrons. Most of the remaining 1% of carbon consists of atoms of the isotope ¹³₆C, with 7 neutrons. A third, even rarer isotope, ¹⁴C, has 8 neutrons. Notice that all three isotopes of carbon have 6 protons; otherwise, they would not be carbon. Although the isotopes of an element have slightly different masses, they behave identically in chemical reactions. (For an element with more than one naturally occurring isotope, the atomic mass is an average of those isotopes, weighted by their abundance. Thus, carbon has an atomic mass of 12.01 daltons.)

Both ¹²C and ¹³C are stable isotopes, meaning that their nuclei do not have a tendency to lose subatomic particles, a process called decay. The isotope ¹⁴C, however, is unstable, or radioactive. A **radioactive isotope** is one in which the nucleus decays spontaneously, giving off particles and energy. When the radioactive decay leads to a change in the number of protons, it transforms the atom to an atom of a different element. For example, when a carbon-14 (¹⁴C) atom decays, a neutron decays into a proton, transforming the atom into a nitrogen (¹⁴N) atom.

Radioactive isotopes have many useful applications in biology. For example, researchers use measurements of radioactivity in fossis to date these relics of past life (see Concept 23.1). Radioactive isotopes are also useful as tracers to follow atoms through metabolism, the chemical processes of an organism. Cells can use radioactive atoms just as they would use nonradioactive isotopes of the same element. The radioactive isotopes are incorporated into biologically active molecules, which can then be tracked by monitoring the radioactivity.

Radioactive tracers are important diagnostic tools in medicine. For example, certain kidney disorders can be diagnosed by injecting small doses of substances containing radioactive isotopes into the blood and then analyzing the tracer molecules excreted in the urine. Radioactive tracers are also used

➤ Figure 2.4 A PET scan, a medical use for radioactive isotopes. PET (positron-emission tomography) detects locations of intense chemical activity in the body. The bright yellow spot marks an area with an elevated level of radioactively labeled glucose, which in turn indicates high metabolic activity, a hallmark of cancerous tissue.



in combination with sophisticated imaging instruments, such as PET scanners, that can monitor the growth and metabolism of cancers in the body (Figure 2.4).

Although radioactive isotopes are very useful in biological research and medicine, radiation from decaying isotopes also poses a hazard to life by damaging cellular molecules. The severity of this damage depends on the type and amount of radiation an organism absorbs. One of the most serious environmental threats is radioactive fallout from nuclear accidents. The doses of most isotopes used in medical diagnosis, however, are relatively safe.

The Energy Levels of Electrons

The simplified models of the atom in Figure 2.3 greatly exaggerate the size of the nucleus relative to that of the whole atom. If an atom of helium were the size of a typical football stadium, the nucleus would be the size of a pencil eraser in the center of the field. Moreover, the electrons would be like two tiny gnats buzzing around the stadium. Atoms are mostly empty space.

When two atoms approach each other during a chemical reaction, their nuclei do not come close enough to interact. Of the three kinds of subatomic particles we have discussed, only electrons are directly involved in the chemical reactions between atoms.

An atom's electrons vary in the amount of energy they possess. **Energy** is defined as the capacity to cause change—for instance, by doing work. **Potential energy** is the energy that matter possesses because of its location or structure. For example, water in a reservoir on a hill has potential energy because of its altitude. When the gates of the reservoir's dam are opened and the water runs downhill, the energy can be used to do work, such as moving the blades of turbines to generate electricity. Because energy has been expended, the water has less energy at the bottom of the hill than it did in the reservoir. Matter has a natural tendency to move toward the lowest possible state of potential energy; in our example, the water runs downhill. To restore the potential energy of a reservoir, work must be done to elevate the water against gravity.


The electrons of an atom have potential energy due to their distance from the nucleus. The negatively charged electrons


are attracted to the positively charged nucleus. It takes work to move a given electron farther away from the nucleus, so the more distant an electron is from the nucleus, the greater its potential energy. Unlike the continuous flow of water downhill, changes in the potential energy of electrons can occur only in steps of fixed amounts. An electron having a certain amount of energy is something like a ball on a staircase (Figure 2.5a). The ball can have different amounts of potential energy, depending on which step it is on, but it cannot spend much time between the steps. Similarly, an electron's potential energy is determined by its energy level. An electron can exist only at certain energy levels, not between them.

An electron's energy level is correlated with its average distance from the nucleus. Electrons are found in different **electron shells**, each with a characteristic average distance and energy level. In diagrams, shells can be represented by concentric circles, as they are in **Figure 2.5b**. The first shell is closest to the nucleus, and electrons in this shell have the lowest potential energy. Electrons in the second shell have more energy, and electrons in the third shell even more energy. An electron can move from one shell to another, but only by absorbing or losing an amount of energy equal to the difference in potential energy between its position in the old shell and that in the new shell. When an electron absorbs energy, it moves to a shell farther out from the nucleus. For example, light energy can excite an electron to a higher energy level. (Indeed, this is the first step

▼ Figure 2.5 Energy levels of an atom's electrons. Electrons exist only at fixed levels of potential energy called electron shells.

(a) A ball bouncing down a flight of stairs can come to rest only on each step, not between steps. Similarly, an electron can exist only at certain energy levels, not between levels.





- (b) An electron can move from one shell to another only if the energy it gains or loses is exactly equal to the difference in energy between the energy levels of the two shells. Arrows in this model indicate some of the stepwise changes in potential energy that are possible.
- Mastering Biology Figure Walkthrough

taken when plants harness the energy of sunlight for photosynthesis, the process that produces food from carbon dioxide and water; see Concept 8.2.) When an electron loses energy, it "falls back" to a shell closer to the nucleus, and the lost energy is usually released to the environment as visible light or ultraviolet radiation.

Electron Distribution and Chemical Properties


The chemical behavior of an atom is determined by the distribution of electrons in the atom's electron shells. Beginning with hydrogen, the simplest atom, we can imagine building the atoms of the other elements by adding 1 proton and 1 electron at a time (along with an appropriate number of neutrons). **Figure 2.6**, a modified version of what is called the *periodic table of the elements*, shows this distribution of electrons for the first 18 elements, from hydrogen ($_1$ H) to argon ($_1$ Ar). The elements are arranged in three rows, or *periods*, corresponding to the number of electron shells in their atoms. The left-to-right sequence of elements in each row corresponds to the sequential addition of electrons and protons. (See Appendix B for the complete periodic table.)

Hydrogen's 1 electron and helium's 2 electrons are located in the first shell. Electrons, like all matter, tend to exist in the lowest available state of potential energy. In an atom, this state is in the first shell. However, the first shell can hold no more than 2 electrons; thus, hydrogen and helium are the only elements in the first row of the table. In an atom with more than 2 electrons, the additional electrons must occupy higher shells because the first shell is full. The next element, lithium, has 3 electrons. Two of these electrons fill the first shell, while the third electron occupies the second shell. The second shell holds a maximum of 8 electrons. Neon, at the end of the second row, has 8 electrons in the second shell, giving it a total of 10 electrons.

The chemical behavior of an atom depends mostly on the number of electrons in its outermost shell. We call those outer electrons valence electrons and the outermost electron shell the valence shell. In the case of lithium, there is only 1 valence electron, and the second shell is the valence shell. Atoms with the same number of electrons in their valence shells exhibit similar chemical behavior. For example, fluorine (F) and chlorine (Cl) both have 7 valence electrons, and both form compounds when combined with the element sodium (Na): Sodium fluoride (NaF) is commonly added to toothpaste to prevent tooth decay, and, as described earlier, NaCl is table salt (see Figure 2.2). An atom with a completed valence shell is unreactive; that is, it will not interact readily with other atoms. At the far right of the periodic table are helium, neon, and argon, the only three elements shown in Figure 2.6 that have full valence shells. These elements are said to be inert, meaning chemically unreactive. All the other atoms in Figure 2.6 are chemically reactive because they have incomplete valence shells.

Notice that as we "build" the atoms in Figure 2.6, the first 4 electrons added to the second and third shells are not shown

▼ Figure 2.6 Electron distribution diagrams for the first 18 elements in the periodic table. In a standard periodic table (see Appendix B), information for each element is presented as shown for helium in the inset. In the diagrams in this table, electrons are represented as yellow dots and electron shells as concentric circles. These diagrams are a convenient way to picture the distribution of an atom's electrons among its electron shells, but these simplified models do not accurately represent the shape of the atom or the location of its electrons. The elements are arranged in rows, each representing the filling of an electron shell. As electrons are added, they occupy the lowest available shell.



VISUAL SKILLS What is the atomic number of magnesium? How many protons and electrons does it have? How many electron shells? How many valence electrons?

in pairs; only after 4 electrons are present do the next electrons complete pairs. The reactivity of an atom arises from the presence of one or more unpaired electrons in its valence shell. As you will see in the next section, atoms interact in a way that completes their valence shells. When they do so, it is the *unpaired* electrons that are involved.

CONCEPT CHECK 2.2

- A nitrogen atom has 7 protons, and the most common isotope of nitrogen has 7 neutrons. A radioactive isotope of nitrogen has 8 neutrons. Write the atomic number and mass number of this radioactive nitrogen as a chemical symbol with a subscript and superscript.
- 2. How many electrons does fluorine have? How many electron shells? How many electrons are needed to fill the valence shell?
- 3. VISUAL SKILLS In Figure 2.6, if two or more elements are in the same row, what do they have in common? If two or more elements are in the same column, what do they have in common?

For suggested answers, see Appendix A.

Mastering Biology Animation: Electron Distribution Diagrams

CONCEPT 2.3

The formation and function of molecules depend on chemical bonding between atoms

Now that we have looked at the structure of atoms, we can move up the hierarchy of organization and see how atoms combine to form molecules and ionic compounds. Atoms with incomplete valence shells can interact with certain other atoms in such a way that each partner completes its valence shell: The atoms either share or transfer valence electrons. These interactions usually result in atoms staying close together, held by attractions called **chemical bonds**. The strongest kinds of chemical bonds are covalent bonds and ionic bonds in dry ionic compounds. (Ionic bonds in aqueous solutions are weak.)

Mastering Biology Animation: Introduction to Chemical Bonds

Covalent Bonds


A **covalent bond** is the sharing of a pair of valence electrons by two atoms. For example, let's consider what happens when two hydrogen atoms approach each other. Recall that hydrogen has 1 valence electron in the first shell, but the shell's capacity is 2 electrons. When the two hydrogen atoms come close enough for their electron shells to overlap, they can share their electrons (**Figure 2.7**). Each hydrogen atom is now associated with 2 electrons in what amounts to a completed valence shell. Two or more atoms held together by covalent bonds constitute a **molecule**, in this case a hydrogen molecule.

Figure 2.8a shows several ways of representing a hydrogen molecule. Its *molecular formula*, H₂, simply indicates that the molecule consists of two atoms of hydrogen. Electron sharing can be depicted by an electron distribution diagram or by a *structural formula*, H—H, where the line represents a **single bond**, a pair of shared electrons. A space-filling model comes closest to representing the actual shape of the molecule.

Oxygen has 6 electrons in its second electron shell and therefore needs 2 more electrons to complete its valence shell. Two oxygen atoms form a molecule by sharing *two* pairs of valence electrons (Figure 2.8b). The atoms are thus joined by a **double bond** (O = O).

Each atom that can share valence electrons has a bonding capacity corresponding to the number of covalent bonds the atom can form. When the bonds form, they give the atom a full complement of electrons in the valence shell. The bonding capacity of oxygen, for example, is 2. This bonding capacity is

▼ Figure 2.7 Formation of a covalent bond.



▼ Figure 2.8 Covalent bonding in four molecules. The number of electrons required to complete an atom's valence shell generally determines how many covalent bonds that atom will form. This figure shows several ways of indicating covalent bonds.

Name and Molecular Formula	Electron Distribution Diagram	Structural Formula	Space- Filling Model
(a) Hydrogen (H ₂). Two hydrogen atoms share one pair of electrons, forming a single bond.	H	н—н	0
(b) Oxygen (O ₂). Two oxygen atoms share two pairs of electrons, forming a double bond.		0=0	
(c) Water (H ₂ O). Two hydrogen atoms and one oxygen atom are joined by single bonds, forming a molecule of water.	H	О—Н Н	9
(d) Methane (CH ₄). Four hydrogen atoms can satisfy the valence of one carbon atom, forming methane.		H H—C—H H	9

Mastering Biology Animation: Covalent Bonds

called the atom's **valence** and usually equals the number of electrons required to complete the atom's outermost (valence) shell. See if you can determine the valences of hydrogen, oxygen, nitrogen, and carbon by studying the electron distribution diagrams in Figure 2.6. You can see that the valence of hydrogen is 1; oxygen, 2; nitrogen, 3; and carbon, 4. (The situation is more complicated for phosphorus, in the third row of the periodic table, which can have a valence of 3 or 5 depending on the combination of single and double bonds it makes.)

The molecules H_2 and O_2 are pure elements rather than compounds because a compound is a combination of two or more *different* elements. Water, with the molecular formula H_2O , is a compound. Two atoms of hydrogen are needed to satisfy the valence of one oxygen atom. **Figure 2.8c** shows the structure of a water molecule. Water is so important to life that the last section of this chapter, Concept 2.5, is devoted to its structure and behavior.

Methane, the main component of natural gas, is a compound with the molecular formula CH₄. It takes four hydrogen atoms, each with a valence of 1, to complete the valence shell of a carbon atom, with its valence of 4 **(Figure 2.8d)**. (We'll look at other carbon compounds in Chapter 3.)


Atoms in a molecule attract shared bonding electrons to varying degrees, depending on the element. The attraction of a particular atom for the electrons of a covalent bond is called its **electronegativity**. The more electronegative an atom is, the more strongly it pulls shared electrons toward itself. In a covalent bond between two atoms of the same element, the electrons are shared equally because the two atoms have the same electronegativity—the tug-of-war is at a standoff. Such a bond is called a **nonpolar covalent bond**. For example, the single bond of H_2 is nonpolar, as is the double bond of O_2 . However, when an atom is bonded to a more electronegative atom, the electrons of the bond are not shared equally. This type of bond is called a polar covalent bond. Such bonds vary in their polarity, depending on the relative electronegativity of the two atoms. For example, the bonds between the oxygen and hydrogen atoms of a water molecule are quite polar (Figure 2.9). Oxygen is one of the most electronegative elements, attracting shared electrons much more strongly than hydrogen does. In a covalent bond between oxygen and hydrogen, the electrons spend more time near the oxygen nucleus than they do near the hydrogen nucleus. Because electrons have a negative charge and are pulled toward oxygen in a water molecule, the oxygen atom has two regions of partial negative charge (each indicated by the Greek letter δ with a minus sign, δ , or "delta minus"), and each hydrogen atom has a partial positive charge (δ +, or "delta plus"). In contrast, the individual bonds of methane (CH₄) are much less polar because the electronegativities of carbon and hydrogen are quite similar.

Ionic Bonds

In some cases, two atoms are so unequal in their attraction for valence electrons that the more electronegative atom strips an electron completely away from its partner. The two resulting oppositely charged atoms (or molecules) are called ions. A positively charged ion is called a cation, while a negatively charged ion is called an anion. Because of their opposite charges, cations and anions attract each other; this attraction is called an ionic bond. Note that the transfer of an electron is not, by itself, the formation of a bond; rather, it allows a bond to form because it results in two ions of opposite charge. Any two such ions can form an ionic bond—the ions do not need to have acquired their charge by an electron transfer with each other.

▼ Figure 2.9 Polar covalent bonds in a water molecule.

Because oxygen (O) is more electronegative than hydrogen (H), shared electrons are pulled more toward oxygen.




Mastering Biology Animation: Nonpolar and Polar Molecules


This is what happens when an atom of sodium ($_{11}$ Na) encounters an atom of chlorine ($_{17}$ Cl) (**Figure 2.10**). A sodium atom has a total of 11 electrons, with its single valence electron in the third electron shell. A chlorine atom has a total of 17 electrons, with 7 electrons in its valence shell. When these two atoms meet, the lone valence electron of sodium is transferred to the chlorine atom, and both atoms end up with their valence shells complete. (Because sodium no longer has an electron in the third shell, the second shell is now the valence shell.)

The electron transfer between the two atoms moves one unit of negative charge from sodium to chlorine. Sodium, now with 11 protons but only 10 electrons, has a net electrical charge of 1+; the sodium atom has become a cation. Conversely, the chlorine atom, having gained an extra electron, now has 17 protons and 18 electrons, giving it a net electrical charge of 1-; it has become a chloride ion—an anion.

▼ Figure 2.10 Electron transfer and ionic bonding. The attraction between oppositely charged atoms, or ions, is an ionic bond. An ionic bond can form between any two oppositely charged ions, even if they have not been formed by transfer of an electron from one to the other.



Mastering Biology Animation: Formation of Ions and Ionic Bonds



▲ Figure 2.11 A sodium chloride (NaCl) crystal. The sodium ions (Na⁺) and chloride ions (Cl⁻) are held together by ionic bonds. The formula NaCl tells us that the ratio of Na⁺ to Cl⁻ is 1:1.

Compounds formed by ionic bonds are called **ionic compounds**, or **salts**. We know the ionic compound sodium chloride (NaCl) as table salt (**Figure 2.11**). Salts are often found in nature as crystals of various sizes and shapes. Each salt crystal is an aggregate of vast numbers of cations and anions bonded by their electrical attraction and arranged in a three-dimensional lattice. Unlike a covalent compound, which consists of molecules having a definite size and number of atoms, an ionic compound does not consist of molecules. The formula for an ionic compound, such as NaCl, indicates only the ratio of elements in a crystal of the salt. "NaCl" by itself is not a molecule.

Not all salts have equal numbers of cations and anions. For example, the ionic compound magnesium chloride (MgCl₂) has two chloride ions for each magnesium ion. Magnesium ($_{12}$ Mg) must lose 2 outer electrons if the atom is to have a complete valence shell, so it has a tendency to become a cation with a net charge of 2+ (Mg²⁺). One magnesium cation can therefore form ionic bonds with two chloride anions (Cl⁻).


The term *ion* also applies to entire molecules that are electrically charged. In the salt ammonium chloride (NH₄Cl), for instance, the anion is a single chloride ion (Cl $^-$), but the cation is ammonium (NH₄ $^+$), a nitrogen atom covalently bonded to four hydrogen atoms. The whole ammonium ion has an electrical charge of 1+ because it has given up 1 electron and thus is 1 electron short.

Environment affects the strength of ionic bonds. In a dry salt crystal, the bonds are so strong that it takes a hammer and chisel to break enough of them to crack the crystal in two. If the same salt crystal is dissolved in water, however, the ionic bonds are much weaker because each ion is partially shielded by its interactions with water molecules. Most drugs are manufactured as salts because they are quite stable when dry but can dissociate (come apart) easily in water.

Weak Chemical Interactions

In organisms, most of the strongest chemical bonds are covalent bonds, which link atoms to form a cell's molecules. But weaker interactions within and between molecules are also indispensable in the cell, contributing greatly to the emergent properties of life. Many large biological molecules are held in their functional form by weak interactions. In addition, when two molecules in the cell make contact, they may adhere temporarily by weak interactions. The reversibility of weak

▼ Figure 2.12 A hydrogen bond.



interactions can be an advantage: Two molecules can come together, affect one another in some way, and then separate.

Several types of weak chemical interactions are important in organisms. One is the ionic bond as it exists between ions dissociated in water, which we just discussed. Hydrogen bonds and van der Waals interactions are also crucial to life.

Hydrogen Bonds

Among the various weak chemical interactions, hydrogen bonds are so central to the chemistry of life that they deserve special attention. When a hydrogen atom is covalently bonded to an electronegative atom, the hydrogen atom has a partial positive charge that allows it to be attracted to a different electronegative atom nearby. This noncovalent attraction between a hydrogen and an electronegative atom is called a **hydrogen bond**. In living cells, the electronegative partners are usually oxygen or nitrogen atoms. Refer to **Figure 2.12** to examine the simple case of hydrogen bonding between water (H_2O) and ammonia (NH_3).

Van der Waals Interactions

Even a molecule with nonpolar covalent bonds may have positively and negatively charged regions. Electrons are not always symmetrically distributed in such a molecule; at any instant, they may accumulate by chance in one part of the molecule or another. The results are ever-changing regions of positive and negative charge that enable all atoms and molecules to stick to one another. These van der Waals interactions are weak and occur only when atoms and molecules are very close together. When many occur simultaneously, however, they can be powerful: They allow a gecko to walk straight up a wall!

A gecko toe has hundreds of thousands of tiny hairs, each with multiple projections, which help to maximize surface contact with the wall. The van der Waals interactions that occur between the molecules of the gecko's foot and those of the wall's surface are so numerous that despite their individual weakness, together they can support the gecko's body weight. This discovery has inspired development of an artificial adhesive called Geckskin: A patch the size of an index card can hold a 700-pound weight to a wall!

Van der Waals interactions, hydrogen bonds, ionic bonds in water, and other weak bonds may form not only between molecules but also between parts of a large molecule, such as a protein. The cumulative effect of weak bonds is to reinforce the three-dimensional shape of the molecule. (You will learn more about the very important biological roles of weak bonds in Concept 3.5.)

Molecular Shape and Function


A molecule has a characteristic size and shape, which are key to its function in the living cell. A molecule consisting of two atoms, such as H_2 or O_2 , is always linear, but most molecules with more than two atoms have more complicated shapes. To take a very simple example, a water molecule (H_2O) is shaped roughly like a V, with its two covalent bonds spread apart at an angle of 104.5° (Figure 2.13). A methane molecule (CH_4) has a geometric shape called a tetrahedron, a pyramid with a triangular base. The carbon nucleus is inside, at the center, with its four covalent bonds radiating to hydrogen nuclei at the corners of the tetrahedron. Larger molecules containing multiple carbon atoms, including many of the molecules that make up living matter, have more complex overall shapes. However, the tetrahedral shape of a carbon atom bonded to four other atoms is often a repeating motif within such molecules.

▼ Figure 2.13 Models showing the shapes of two small molecules. Each of the molecules, water and methane, is represented in two different ways.


Name and Molecular Formula	Space-Filling Model	Ball-and-Stick Model
Water (H ₂ O)		H 104.5° H
Methane (CH4)		H

Molecular shape is crucial in biology: It determines how biological molecules recognize and respond to one another with specificity. Biological molecules often bind temporarily to each other by forming weak interactions, but only if their shapes are complementary. Consider the effects of opiates (drugs derived from opium); morphine and heroin are two examples. Opiates relieve pain and alter mood by weakly binding to specific receptor molecules on the surfaces of brain cells. Why would brain cells carry receptors for opiates, compounds that are not endogenous, made by the body? In 1975, this question was answered by the discovery of endorphins ("endogenous morphines"). Endorphins are signaling molecules made by the pituitary gland that bind to the receptors, relieving pain and producing euphoria during times of stress, such as intense exercise. Opiates have shapes similar to endorphins and mimic them by binding to endorphin receptors in the brain. That is why opiates and endorphins have similar effects (Figure 2.14).

▼ Figure 2.14 A molecular mimic. Morphine affects pain perception and emotional state by mimicking the brain's natural endorphins.



(a) Structures of endorphin and morphine. The boxed portion of the endorphin molecule (left) binds to receptor molecules on target cells in the brain. The boxed portion of the morphine molecule (right) is a close match.



- (b) Binding to endorphin receptors. Both endorphin and morphine can bind to endorphin receptors on the surface of a brain cell.
- Mastering Biology Interview with Candace Pert: Discovering opiate receptors in the brain



The role of molecular shape in brain chemistry illustrates the match between structure and function in biological organization, one of biology's unifying themes.


CONCEPT CHECK 2.3

- 1. Why does the structure H—C—C—H fail to make sense chemically?
- 2. What holds the atoms together in a crystal of magnesium chloride (MgCl₂)?
- 3. WHAT IF? If you were a pharmaceutical researcher, why would you want to learn the three-dimensional shapes of naturally occurring signaling molecules?
 For suggested answers, see Appendix A.

CONCEPT 2.4

Chemical reactions make and break chemical bonds


The making and breaking of chemical bonds, leading to changes in the composition of matter, are called **chemical reactions**. An example is the reaction between hydrogen and oxygen molecules that forms water:



This reaction breaks the covalent bonds of H_2 and O_2 and forms the new bonds of H_2O . When we write the equation for a chemical reaction, we use an arrow to indicate the conversion of the starting materials, called the **reactants**, to the resulting materials, or **products**. The coefficients indicate the number of molecules involved; for example, the coefficient 2 in front of H_2 means that the reaction starts with two molecules of hydrogen. Notice that all atoms of the reactants must be accounted for in the products. Matter is conserved in a chemical reaction: Reactions cannot create or destroy atoms but can only rearrange (redistribute) the electrons among them.

Photosynthesis, which takes place within the cells of green plant tissues, is an important biological example of how chemical reactions rearrange matter. Humans and other animals ultimately depend on photosynthesis for food and oxygen, and this process is at the foundation of life in almost all ecosystems.

The following diagram summarizes photosynthesis:



The raw materials of photosynthesis are carbon dioxide (CO_2) and water (H_2O) , which land plants absorb from the air and soil, respectively. Within the plant cells, sunlight powers the conversion of these ingredients to a sugar called glucose $(C_6H_{12}O_6)$ and oxygen molecules (O_2) , a by-product that can be seen when released by a water plant **(Figure 2.15)**. Although photosynthesis is actually a sequence of many chemical reactions, we still end up with the same number and types of atoms that we had when we started. Matter has simply been rearranged, with an input of energy provided by sunlight.

All chemical reactions are theoretically reversible, with the products of the forward reaction becoming the reactants for the reverse reaction. For example, hydrogen and nitrogen molecules can combine to form ammonia, but ammonia can also decompose to regenerate hydrogen and nitrogen:

$$3 H_2 + N_2 \rightleftharpoons 2 NH_3$$

The two opposite-headed arrows indicate that the reaction is reversible.

One of the factors affecting the rate of a reaction is the concentration of reactants. The greater the concentration of reactant molecules, the more frequently they collide with one



▼Figure 2.15 Photosynthesis: a solar-powered rearrangement of matter. Elodea, a freshwater plant, produces sugar by rearranging the atoms of carbon dioxide and water in the chemical process known as photosynthesis, which is powered by sunlight. Much of the sugar is then converted to other food molecules. Oxygen gas (O₂) is a byproduct of photosynthesis; notice the bubbles of O₂-containing gas escaping from the leaves submerged in water.

VISUAL SKILLS Add labels and arrows on the photo showing the reactants and products of photosynthesis as it takes place in a leaf.

another and have an opportunity to react and form products. The same holds true for products. As products accumulate, collisions resulting in the reverse reaction become more frequent. Eventually, the forward and reverse reactions occur at the same rate, and the relative concentrations of products and reactants stop changing. The point at which the reactions offset one another exactly is called chemical equilibrium. This is a dynamic equilibrium; reactions are still going on in both directions, but with no net effect on the concentrations of reactants and products. Equilibrium does not mean that the reactants and products are equal in concentration, but only that their concentrations have stabilized at a particular ratio. The reaction involving ammonia reaches equilibrium when ammonia decomposes as rapidly as it forms. In some chemical reactions, the equilibrium point may lie so far to the right that these reactions go essentially to completion; that is, virtually all the reactants are converted to products.

To conclude this chapter, we focus on water, the substance in which all the chemical processes of organisms occur.

CONCEPT CHECK 2.4

- 1. Which type of chemical reaction, if any, occurs faster at equilibrium, the formation of products from reactants or that of reactants from products?
- WHAT IF? Write an equation that uses the products of photosynthesis as reactants and the reactants of photosynthesis as products. Add energy as another product. This new equation describes a process that occurs in your cells. Describe this equation in words. How does this equation relate to breathing?

For suggested answers, see Appendix A.

CONCEPT 2.5

Hydrogen bonding gives water properties that help make life possible on Earth

All organisms are made mostly of water and live in an environment dominated by water. Most cells are surrounded by water, and cells themselves are about 70–95% water. Water is so familiar to us that it is easy to overlook its extraordinary qualities. We can trace water's unique behavior to the structure and interactions of its molecules. As you saw in Figure 2.9, the connections between the atoms of a water molecule are polar covalent bonds. The unequal sharing of electrons and water's V-like shape make it a **polar molecule**, meaning that its overall charge is unevenly distributed: The oxygen region of the molecule has two partial negative charges $(\delta-)$, and each hydrogen has a partial positive charge $(\delta+)$.

The properties of water arise from attractions between oppositely charged atoms of different water molecules: The partially


positive hydrogen of one molecule is attracted to the partially negative oxygen of a nearby molecule. The two molecules are thus held together by a hydrogen bond. When water is in its liquid form, its hydrogen bonds are very fragile, each only about 1/20 as strong as a covalent bond. The hydrogen bonds form, break, and re-form with great frequency. Each lasts only a few trillionths of a second, but the molecules are constantly forming new hydrogen bonds with a succession of partners. Therefore, at any instant, most of the water molecules are hydrogen-bonded to their neighbors (Figure 2.16). The extraordinary properties of water emerge from this hydrogen bonding, which organizes water molecules into a higher level of structural order. We will examine four emergent properties of water that contribute to Earth's suitability as an environment for life: cohesive behavior, ability to moderate temperature, expansion upon freezing, and versatility as a solvent. After that, we'll discuss a critical aspect of water chemistry—acids and bases.

Cohesion of Water Molecules

Water molecules stay close to each other as a result of hydrogen bonding. At any given moment, many of the molecules in liquid water are linked by multiple hydrogen bonds. These linkages make water more structured than most other liquids. Collectively, the hydrogen bonds hold the substance together, a phenomenon called **cohesion**.

▼ Figure 2.16 Hydrogen bonds between water molecules.

The charged regions in a water molecule are due to its polar covalent bonds. Oppositely charged regions of neighboring water molecules are attracted to each other, forming hydrogen bonds. Each molecule can hydrogen-bond to multiple partners, and these associations are constantly changing.




DRAW IT Draw partial charges on all the atoms of the water molecule on the far left, and draw three more water molecules hydrogen-bonded to it.

Mastering Biology Animation: Polarity of Water Cohesion due to hydrogen bonding contributes to the transport of water and dissolved nutrients against gravity in plants (Figure 2.17). Water from the roots reaches the leaves through a network of water-conducting cells. As water evaporates from a leaf, hydrogen bonds cause water molecules leaving the veins to tug on molecules farther down, and the upward pull is transmitted through the water-conducting cells all the way to the roots. Adhesion, the clinging of one substance to another, also plays a role. Adhesion of water to cell walls by hydrogen bonds helps counter the downward pull of gravity.

Related to cohesion is **surface tension**, a measure of how difficult it is to stretch or break the surface of a liquid. At the air-water interface is an ordered arrangement of water molecules, hydrogen bonded to one another and to the water below, but not to the air above. This asymmetry gives water an unusually high surface tension, making it behave as though it were coated with an invisible film. You can observe the surface tension of water by slightly overfilling a drinking glass; the water will stand above the rim. The spider in **Figure 2.18** takes advantage of the surface tension of water to walk across a pond without breaking the surface.

▼ Figure 2.17 Water transport in plants. Evaporation from leaves pulls water upward from the roots through water-conducting cells. Because of the properties of cohesion and adhesion, the tallest trees can transport water more than 100 m upward—approximately one-quarter the height of the Empire State Building in New York City.



 Mastering Biology BioFlix[®] Animation: Adhesion and Cohesion in Plants Animation: Cohesion of Water

Figure 2.18 Walking on water. The high surface tension of water, resulting from the collective strength of its hydrogen bonds, allows this raft spider to walk on the surface of a pond.

Moderation of Temperature by Water

Water moderates air temperature by absorbing heat from air that is warmer and releasing the stored heat to air that is cooler. Water is effective as a heat bank because it can absorb or release a relatively large amount of heat with only a slight change in its own temperature. To understand this capability of water, let's first look at temperature and heat.

Temperature and Heat

Anything that moves has kinetic energy, the energy of motion. Atoms and molecules have kinetic energy because they are always moving, although not necessarily in any particular direction. The faster a molecule moves, the greater its kinetic energy. The kinetic energy associated with the random movement of atoms or molecules is called thermal energy. Thermal energy is related to temperature, but they are not the same thing. Temperature represents the average kinetic energy of the molecules in a body of matter, regardless of volume, whereas the thermal energy of a body of matter reflects the total kinetic energy and thus depends on the matter's volume. When water is heated in a coffeemaker, the average speed of the molecules increases, and the thermometer records this as a rise in temperature of the liquid. The total amount of thermal energy also increases in this case. Note, however, that although the pot of coffee has a much higher temperature than, say, the water in a swimming pool, the swimming pool contains more thermal energy because of its much greater volume.

Whenever two objects of different temperature are brought together, thermal energy passes from the warmer to the cooler object until the two are the same temperature. Molecules in the cooler object speed up at the expense of the thermal energy of the warmer object. An ice cube cools a drink not by adding coldness to the liquid but by absorbing thermal energy from the liquid as the ice itself melts. Thermal energy in transfer from one body of matter to another is defined as **heat**.

One convenient unit of heat used in this book is the **calorie (cal)**. A calorie is the amount of heat it takes to raise the temperature of 1 g of water by 1°C. Conversely, a calorie is

also the amount of heat that 1 g of water releases when it cools by 1°C. A **kilocalorie** (**kcal**), 1,000 cal, is the quantity of heat required to raise the temperature of 1 kilogram (kg) of water by 1°C. (The "Calories" on food packages are actually kilocalories.) Another energy unit used in this book is the **joule** (**J**). One joule equals 0.239 cal; one calorie equals 4.184 J.

Water's High Specific Heat

The ability of water to stabilize temperature stems from its relatively high specific heat. The specific heat of a substance is defined as the amount of heat that must be absorbed or lost for 1 g of that substance to change its temperature by 1°C. We already know water's specific heat because we have defined a calorie as the amount of heat that causes 1 g of water to change its temperature by 1°C. Therefore, the specific heat of water is 1 calorie per gram per degree Celsius, abbreviated as 1 cal/(g · °C). Compared with most other substances, water has an unusually high specific heat. As a result, water will change its temperature less than other liquids when it absorbs or loses a given amount of heat. The reason you can burn your fingers by touching the side of an iron pot on the stove when the water in the pot is still lukewarm is that the specific heat of water is ten times greater than that of iron. In other words, the same amount of heat will raise the temperature of 1 g of the iron much faster than it will raise the temperature of 1 g of the water. Specific heat can be thought of as a measure of how well a substance resists changing its temperature when it absorbs or releases heat. Water resists changing its temperature; when it does change its temperature, it absorbs or loses a relatively large quantity of heat for each degree of change.

We can trace water's high specific heat, like many of its other properties, to hydrogen bonding. Heat must be absorbed in order to break hydrogen bonds; by the same token, heat is released when hydrogen bonds form. A calorie of heat causes a relatively small change in the temperature of water because much of the heat is used to disrupt hydrogen bonds before the water molecules can begin moving faster. And when the temperature of water drops slightly, many additional hydrogen bonds form, releasing a considerable amount of energy in the form of heat.

What is the relevance of water's high specific heat to life on Earth? A large body of water can absorb and store a huge amount of heat from the sun in the daytime and during summer while warming up only a few degrees. At night and during winter, the gradually cooling water can warm the air. This capability of water serves to moderate air temperatures in coastal areas (Figure 2.19). The high specific heat of water also tends to stabilize ocean temperatures, creating a favorable environment for marine life. Thus, because of its high specific heat, the water that covers most of Earth keeps temperature fluctuations on land and in water within limits that permit life. Also, because organisms are made primarily of water, they are better able to resist changes in their own temperature than if they were made of a liquid with a lower specific heat.

▼ Figure 2.19 Temperatures for the Pacific Ocean and Southern California on an August day.



INTERPRET THE DATA Explain the pattern of temperatures shown in this diagram.

Evaporative Cooling

Molecules of any liquid stay close together because they are attracted to one another. Molecules moving fast enough to overcome these attractions can depart the liquid and enter the air as a gas (vapor). This transformation from a liquid to a gas is called vaporization, or *evaporation*. Recall that the speed of molecular movement varies and that temperature is the *average* kinetic energy of molecules. Even at low temperatures, the speediest molecules can escape into the air. Some evaporation occurs at any temperature; a glass of water at room temperature, for example, will eventually evaporate completely. If a liquid is heated, the average kinetic energy of molecules increases and the liquid evaporates more rapidly.

Heat of vaporization is the quantity of heat a liquid must absorb for 1 g of it to be converted from the liquid to the gaseous state. For the same reason that water has a high specific heat, it also has a high heat of vaporization relative to most other liquids. To evaporate 1 g of water at 25°C, about 580 cal of heat is needed—nearly double the amount needed to vaporize a gram of alcohol, for example. Water's high heat of vaporization is another emergent property resulting from the strength of its hydrogen bonds, which must be broken before the molecules can exit from the liquid in the form of water vapor.

The high amount of energy required to vaporize water has a wide range of effects. On a global scale, for example, it helps moderate Earth's climate. A considerable amount of solar heat absorbed by tropical seas is consumed during the evaporation of surface water. Then, as moist tropical air circulates poleward, it releases heat as it condenses and forms rain. On an organismal level, water's high heat of vaporization accounts for the severity of steam burns. These burns are caused by the heat energy released (during formation of hydrogen bonds) when steam condenses into liquid on the skin.

As a liquid evaporates, the surface of the liquid that remains behind cools down (its temperature decreases). This **evaporative cooling** occurs because the "hottest" molecules, those with the greatest kinetic energy, are the ones most likely to leave as gas. It is as if the 100 fastest runners at a college transferred to another school; the average speed of the remaining students would decline.

Evaporative cooling of water contributes to the stability of temperature in lakes and ponds and also provides a mechanism that prevents terrestrial organisms from overheating. For example, evaporation of water from the leaves of a plant helps keep the tissues in the leaves from becoming too warm in the sunlight. Evaporation of sweat from human skin dissipates body heat and helps prevent overheating on a hot day or when excess heat is generated by strenuous activity. High humidity on a hot day increases discomfort because the high concentration of water vapor in the air inhibits the evaporation of sweat from the body.

Floating of Ice on Liquid Water

Water is one of the few substances that are less dense as a solid than as a liquid. In other words, ice floats on liquid water. While other materials contract and become denser when they solidify, water expands. The cause of this exotic behavior is, once again, hydrogen bonding. At temperatures above 4°C, water behaves like other liquids, expanding as it warms and contracting as it cools. As the temperature falls from 4°C to 0°C, water begins to freeze because more and more of its molecules are moving too slowly to break hydrogen bonds. At 0°C, the molecules become locked into a crystalline lattice, each water molecule hydrogenbonded to four partners (Figure 2.20). The hydrogen bonds keep the molecules at "arm's length," far enough apart to make ice about 10% less dense than liquid water at 4°C. When ice absorbs enough heat for its temperature to rise above 0°C, hydrogen bonds between molecules are disrupted. As the crystal collapses, the ice melts, and molecules have fewer hydrogen bonds, allowing them to slip closer together. Water reaches its greatest density at 4°C and then begins to expand as the molecules move faster.

The ability of ice to float due to its lower density is an important factor in the suitability of the environment for life. If ice sank, then eventually all ponds, lakes, and even oceans would

freeze solid, making life as we know it impossible on Earth. During summer, only the upper summer, only the upper summer, only the upper summer. Hydrogen bonds break and re-form

Ice:
Hydrogen bonds are stable

few inches of the ocean would thaw. Instead, when a deep body of water cools, the floating ice insulates the liquid water below, preventing it from freezing and allowing life to exist under the frozen surface, as shown in the photo in Figure 2.20. Besides insulating the water below, ice also provides a solid habitat for some animals, such as polar bears, seals, and walruses.

Many scientists are worried that these bodies of ice are at risk of disappearing. Global warming, which is caused by carbon dioxide and other "greenhouse" gases in the atmosphere (see Figure 43.26), is having a profound effect on icy environments around the globe. In the Arctic, the average air temperature has risen 2.2°C just since 1961. This temperature increase has affected the seasonal balance between Arctic sea ice and liquid water, causing ice to form later in the year, to melt earlier, and to cover a smaller area. The rate at which glaciers and Arctic sea ice are disappearing is posing an extreme challenge to animals that depend on ice for their survival (Figure 2.21).

Water: The Solvent of Life

A sugar cube placed in a glass of water will dissolve. In time, the glass will contain a uniform mixture of sugar and water; the concentration of dissolved sugar will be the same everywhere in the mixture. A liquid that is a completely homogeneous mixture of two or more substances is called a **solution**. The dissolving agent of a solution is the **solvent**, and the substance that is dissolved is the **solute**. In this case, water is the solvent and sugar is the solute. An **aqueous solution** is one in which the solute is dissolved in water; water is the solvent.

Water is a very versatile solvent, a quality we can trace to the polarity of the water molecule. Suppose, for example, that a spoonful of table salt, the ionic compound sodium chloride (NaCl), is placed in water (Figure 2.22). At the surface of each grain, or crystal, of salt, the sodium and chloride ions are exposed to the solvent. These ions and regions of the water molecules are attracted to each other due to their opposite charges. The oxygens of the water molecules have regions of partial negative charge that are attracted to sodium cations. The hydrogen regions are

■ Figure 2.20 Ice: crystal structure and floating barrier. In ice, each molecule is hydrogen-bonded to four neighbors in a spacious, three-dimensional crystal. Ice has fewer molecules than an equal volume of liquid water, so ice is less dense than liquid water. Floating ice is a barrier that insulates the liquid water below from the colder air. The marine organism shown here is a type of shrimp called krill; it was photographed beneath floating ice in the Southern Ocean near Antarctica.

WHAT IF? If water did not form hydrogen bonds, what would happen to the shrimp's habitat, shown here?

▶ Figure 2.21 Effects of climate change on the Arctic. Warmer temperatures in the Arctic are causing more sea ice to melt in the summer; the ice sheet reaches its minimum in September. The loss of ice benefits some organisms and harms others. Map data from National Snow and Ice Data Center (http://nsidc.org/arcticseaicenews/2018/10).


Mastering Biology
Interview with Susan Solomon:
Understanding climate change





partially positively charged and are attracted to chloride anions. As a result, water molecules surround the individual sodium and chloride ions, separating and shielding them from one another. The sphere of water molecules around each dissolved ion is called a **hydration shell**. Working inward from the surface of each salt crystal, water eventually dissolves all the ions. The result is a solution of two solutes, sodium cations and chloride anions, mixed homogeneously with water, the solvent. Other ionic compounds also dissolve in water. Seawater, for instance, contains a great variety of dissolved ions, as do living cells.

▼ Figure 2.22 Table salt dissolving in water. A sphere of water molecules, called a hydration shell, surrounds each solute ion.




A compound does not need to be ionic to dissolve in water; many compounds made up of nonionic polar molecules, such as sugars, are also water-soluble. Such compounds dissolve when water molecules surround each of the solute molecules, forming hydrogen bonds with them. Even molecules as large as proteins can dissolve in water if they have ionic and polar regions on their surface (Figure 2.23). Many different kinds of polar compounds are dissolved (along with ions) in the water of such biological fluids as blood, the sap of plants, and the liquid within all cells. Water is the solvent of life.

▼ Figure 2.23 A water-soluble protein. Human lysozyme is a protein found in tears and saliva that has antibacterial action (see Figure 3.20). This model shows the lysozyme molecule (purple) in an aqueous environment. Ionic and polar regions on the protein's surface attract the partially charged

regions on water molecules.

This oxygen is attracted to a slight positive charge on the lysozyme molecule.



This hydrogen is attracted to a slight negative charge on the lysozyme molecule.

Hydrophilic and Hydrophobic Substances

Any substance that has an affinity for water is said to be **hydrophilic** (from the Greek *hydro*, water, and *philos*, loving). In some cases, substances can be hydrophilic without actually dissolving. For example, some molecules in cells are so large that they do not dissolve. Another example of a hydrophilic substance that does not dissolve is cotton, a plant product. Cotton consists of giant molecules of cellulose, a compound with numerous regions of partial positive and partial negative charges that can form hydrogen bonds with water. Water adheres to the cellulose fibers. Thus, a cotton towel does a great job of drying the body, yet it does not dissolve in the washing machine. Cellulose is also present in the walls of plant cells that conduct water; you read earlier how the adhesion of water to these hydrophilic walls helps water move up the plant against gravity.

There are, of course, substances that do not have an affinity for water. Substances that are nonionic and nonpolar (or otherwise cannot form hydrogen bonds) actually seem to repel water; these substances are said to be **hydrophobic** (from the Greek *phobos*, fearing). An example from the kitchen is vegetable oil, which, as you know, does not mix stably with water-based substances such as vinegar. The hydrophobic behavior of the oil molecules results from a high number of relatively nonpolar covalent bonds, in this case bonds between carbon and hydrogen, which share electrons almost equally. Hydrophobic molecules related to oils are major ingredients of cell membranes. (Imagine what would happen to a cell if its membrane dissolved!)

Solute Concentration in Aqueous Solutions

Most of the chemical reactions in organisms involve solutes dissolved in water. To understand such reactions, we must know how many atoms and molecules are involved and calculate the concentration of solutes in an aqueous solution (the number of solute molecules in a volume of solution).

When carrying out experiments, we use mass to calculate the number of molecules. We first calculate the molecular mass, which is simply the sum of the masses of all the atoms in a molecule. As an example, let's calculate the molecular mass of table sugar (sucrose), C12H22O11, by multiplying the number of atoms by the atomic mass of each element (see Appendix B). In round numbers, sucrose has a molecular mass of $(12 \times 12) + (22 \times 1) + (11 \times 16) = 342$ daltons. Because we can't weigh out small numbers of molecules, we usually measure substances in units called moles. Just as a dozen always means 12 objects, a mole (mol) represents an exact number of objects: 6.02×10^{23} , which is called Avogadro's number. There are 6.02×10^{23} daltons in 1 g. Once we determine the molecular mass of a molecule such as sucrose, we can use the same number (342), but with the unit grams, to represent the mass of 6.02×10^{23}


molecules of sucrose, or 1 mol of sucrose (sometimes called the molar mass). To obtain 1 mol of sucrose in the lab, therefore, we weigh out 342 g.

The practical advantage of measuring a quantity of chemicals in moles is that a mole of one substance has exactly the same number of molecules as a mole of any other substance. Measuring in moles makes it convenient for scientists working in the laboratory to combine substances in fixed ratios of molecules.

How would we make a liter (L) of solution consisting of 1 mol of sucrose dissolved in water? We would measure out 342 g of sucrose and then add enough water to bring the total volume of the solution up to 1 L. At that point, we would have a 1-molar (1 *M*) solution of sucrose. **Molarity**—the number of moles of solute per liter of solution—is the unit of concentration most often used by biologists for aqueous solutions.

Acids and Bases

Occasionally, a hydrogen atom participating in a hydrogen bond between two water molecules shifts from one molecule to the other. When this happens, the hydrogen atom leaves its electron behind, and what is actually transferred is a **hydrogen ion** (H^+), a single proton with a charge of 1+. The water molecule that lost a proton is now a **hydroxide ion** (OH^-), which has a charge of 1-. The proton binds to the other water molecule, making that molecule a **hydronium ion** (H_3O^+):



By convention, H^+ (the hydrogen ion) is used to represent H_3O^+ (the hydronium ion), and we follow that practice here. Keep in mind, though, that H^+ does not exist on its own in an aqueous solution. It is always associated with a water molecule in the form of H_3O^+ .

As indicated by the double arrows, this is a reversible reaction that reaches a state of dynamic equilibrium when water molecules dissociate at the same rate that they are being reformed from H^+ and OH^- . At this equilibrium point, the concentration of water molecules greatly exceeds the concentrations of H^+ and OH^- . In pure water, only one water molecule in every 554 million is dissociated; the concentration of H^+ and of OH^- in pure water is therefore $10^{-7}M$ (at $25^{\circ}C$). This means there is only one ten-millionth of a mole of hydrogen ions per liter of pure water and an equal number of hydroxide ions. (Even so, this is a huge number—over 60,000 trillion—of each ion per liter.)

Although the dissociation of water is reversible and statistically rare, it is exceedingly important in the chemistry of life. H^+ and OH^- are very reactive. Changes in their concentrations can drastically affect a cell's proteins and other complex molecules. As we have seen, the concentrations of H^+ and OH^- are equal in pure water, but adding certain kinds of solutes, called acids and bases, disrupts this balance.

What would cause an aqueous solution to have an imbalance in H⁺ and OH⁻ concentrations? When acids dissolve in water, they donate additional H⁺ to the solution. An **acid** is a substance that increases the hydrogen ion concentration of a solution. For example, when hydrochloric acid (HCl) is added to water, hydrogen ions dissociate from chloride ions:

This source of H⁺ (dissociation of water is the other source) results in an acidic solution—one having more H⁺ than OH⁻.

A substance that *reduces* the hydrogen ion concentration of a solution is called a **base**. Some bases reduce the H^+ concentration directly by accepting hydrogen ions. Ammonia (NH₃), for instance, acts as a base when the unshared electron pair in nitrogen's valence shell attracts a hydrogen ion from the solution, resulting in an ammonium ion (NH₄⁺):

$$NH_3 + H^+ \Longrightarrow NH_4^+$$

Other bases reduce the H⁺ concentration indirectly by dissociating to form hydroxide ions, which combine with hydrogen ions and form water. One such base is sodium hydroxide (NaOH), which in water dissociates into its ions:

In either case, the base reduces the H^+ concentration. Solutions with a higher concentration of OH^- than H^+ are known as basic solutions. A solution in which the H^+ and OH^- concentrations are equal is said to be neutral.

Notice that single arrows were used in the reactions for HCl and NaOH. These compounds dissociate completely when mixed with water, so hydrochloric acid is called a strong acid and sodium hydroxide a strong base. In contrast, ammonia is a weak base. The double arrows in the reaction for ammonia indicate that the binding and release of hydrogen ions are reversible reactions, although at equilibrium there will be a fixed ratio of NH_4^+ to NH_3 .

Weak acids are acids that reversibly release and accept back hydrogen ions. An example is carbonic acid:

$$H_2CO_3 \Longrightarrow HCO_3^- + H^+$$
Carbonic Bicarbonate Hydrogen
acid ion ion

Here the equilibrium so favors the reaction in the left direction that when carbonic acid is added to pure water, only 1% of the molecules are dissociated at any particular time. Still, that is enough to shift the balance of H⁺ and OH⁻ from neutrality.

The pH Scale

In any aqueous solution at 25°C, the *product* of the H^+ and OH^- concentrations is constant at 10^{-14} . This can be written

$$[H^+][OH^-] = 10^{-14}$$

(The brackets indicate molar concentration.) As previously mentioned, in a neutral solution at room temperature (25°C), $[H^+] = 10^{-7}$ and $[OH^-] = 10^{-7}$. Therefore, the product of [H⁺] and [OH⁻] in a neutral solution at 25°C is 10⁻¹⁴. If enough acid is added to a solution to increase $[H^+]$ to $10^{-5} M$, then [OH-] will decline by an equivalent factor to 10-9 M (note that $10^{-5} \times 10^{-9} = 10^{-14}$). This constant relationship expresses the behavior of acids and bases in an aqueous solution. An acid not only adds hydrogen ions to a solution, but also removes hydroxide ions because of the tendency for H+ to combine with OH⁻, forming water. A base has the opposite effect, increasing OH⁻ concentration but also reducing H⁺ concentration by the formation of water. If enough of a base is added to raise the OH⁻ concentration to 10⁻⁴ M, it will cause the H $^+$ concentration to drop to 10^{-10} M. Whenever we know the concentration of either H⁺ or OH⁻ in an aqueous solution, we can deduce the concentration of the other ion.

Because the H⁺ and OH⁻ concentrations of solutions can vary by a factor of 100 trillion or more, scientists have developed a way to express this variation more conveniently than in moles per liter. The pH scale (Figure 2.24) compresses the range of H⁺ and OH⁻ concentrations by employing logarithms. The **pH** of a solution is defined as the negative logarithm (base 10) of the hydrogen ion concentration:

$$pH = -log[H^+]$$

For a neutral aqueous solution, $[H^+]$ is 10^{-7} M, giving us


$$-\log 10^{-7} = -(-7) = 7$$

Notice that pH *decreases* as H $^+$ concentration *increases* (see Figure 2.24). Notice, too, that although the pH scale is based on H $^+$ concentration, it also implies OH $^-$ concentration. A solution of pH 10 has a hydrogen ion concentration of $10^{-10}\,M$ and a hydroxide ion concentration of $10^{-4}\,M$.

The pH of a neutral aqueous solution at 25°C is 7, the midpoint of the pH scale. A pH value less than 7 denotes an acidic solution; the lower the number, the more acidic the solution. The pH for basic solutions is above 7. Most biological fluids, such as blood and saliva, are within the range of pH 6–8. There are a few exceptions, however, including the strongly acidic digestive juice of the human stomach (gastric juice), which has a pH of about 2.

Remember that each pH unit represents a tenfold difference in H $^+$ and OH $^-$ concentrations. It is this mathematical feature that makes the pH scale so compact. A solution of pH 3 is not twice as acidic as a solution of pH 6, but 1,000 times (10 \times 10 \times 10) more acidic. When the pH of a solution changes slightly, the actual concentrations of H $^+$ and OH $^-$ in the solution change substantially.

▼ Figure 2.24 The pH scale and pH values of some aqueous solutions.



Buffers

The internal pH of most living cells is close to 7. Even a slight change in pH can be harmful because the chemical processes of the cell are very sensitive to the concentrations of hydrogen and hydroxide ions. The pH of human blood is very close to 7.4, which is slightly basic. A person cannot survive for more than a few minutes if the blood pH drops to 7 or rises to 7.8, and a chemical system exists in the blood that maintains a stable pH. If 0.01 mol of a strong acid is added to a liter of pure water, the pH drops from 7.0 to 2.0. If the same amount of acid is added to a liter of blood, however, the pH decrease is only from 7.4 to 7.3. Why does the addition of acid have so much less of an effect on the pH of blood than it does on the pH of water?

The presence of substances called buffers allows biological fluids to maintain a relatively constant pH despite the

addition of acids or bases. A **buffer** is a substance that minimizes changes in the concentrations of H⁺ and OH⁻ in a solution. It does so by accepting hydrogen ions from the solution when they are in excess and donating hydrogen ions to the solution when they have been depleted. Most buffer solutions contain a weak acid and its corresponding base, which combine reversibly with hydrogen ions.

Several buffers contribute to pH stability in human blood and many other biological solutions. One of these is carbonic acid (H_2CO_3), which is formed when CO_2 reacts with water in blood plasma. As mentioned earlier, carbonic acid dissociates to yield a bicarbonate ion (HCO_3^-) and a hydrogen ion (H^+):

The chemical equilibrium between carbonic

acid and bicarbonate acts as a pH regulator, the reaction shifting left or right as other processes in the solution add or remove hydrogen ions. If the H⁺ concentration in blood begins to fall (that is, if pH rises), the reaction proceeds to the right and more carbonic acid dissociates, replenishing hydrogen ions. But when the H⁺ concentration in blood begins to rise (when pH drops), the reaction proceeds to the left, with (HCO₃⁻) (the base) removing the hydrogen ions from the solution and forming H₂CO₃. Thus, the carbonic acid–bicarbonate buffering system consists of an acid and a base in equilibrium with each other. Most other buffers are also acid-base pairs.


Acidification: A Threat to Our Oceans

Among the many threats to water quality posed by human activities is the burning of fossil fuels, which releases CO_2 into the atmosphere. The resulting increase in atmospheric CO_2 levels has caused global warming and other aspects of climate change (see Concept 43.4). In addition, about 25% of human-generated CO_2 is absorbed by the oceans. In spite of the huge volume of water in the oceans, scientists worry that the absorption of so much CO_2 will harm marine ecosystems.

Recent data have shown that such fears are well founded. When CO_2 dissolves in seawater, it reacts with water to form carbonic acid, which lowers ocean pH. This process, known as **ocean acidification**, alters the delicate balance of conditions for life in the oceans (**Figure 2.25**). Based on measurements of CO_2 levels in air bubbles trapped in ice over thousands of years, scientists calculate that the pH of the oceans is 0.1 pH unit lower now than at any time in the past 420,000 years. Recent studies predict that it will drop another 0.3–0.5 pH unit by the end of this century.

As seawater acidifies, the extra hydrogen ions combine with carbonate ions (CO_3^{2-}) to form bicarbonate ions (HCO_3^{-}), thereby reducing the carbonate ion concentration (see Figure 2.25).

▼ Figure 2.25 Ocean acidification.



Some carbon dioxide (CO₂) in the atmosphere dissolves in the ocean, where it reacts with water to form carbonic acid (H₂CO₃).

Carbonic acid dissociates into hydrogen ions (H⁺) and bicarbonate ions (HCO₃⁻).

The added H⁺ combines with carbonate ions (CO₃²⁻), forming more HCO₃⁻.

Less CO₃²⁻ is available for calcification — the formation of calcium carbonate (CaCO₃)— by marine organisms such as corals.

VISUAL SKILLS Summarize the effect of adding excess CO_2 to the oceans on the calcification process in the final equation.

Scientists predict that ocean acidification will cause the carbonate ion concentration to decrease by 40% by the year 2100. This is of great concern because carbonate ions are required for calcification, the production of calcium carbonate (CaCO₃), by many marine organisms, including reef-building corals and animals that build shells. The **Scientific Skills Exercise** allows you to work with data from an experiment examining the effect of carbonate ion concentration on coral reefs. Coral reefs are sensitive ecosystems that act as havens for a great diversity of marine life. The disappearance of coral reef ecosystems would be a tragic loss of biological diversity.

CONCEPT CHECK 2.5

- 1. Describe how properties of water contribute to the upward movement of water in a tree.
- 2. How can the freezing of water crack boulders?
- The same volume of a solution at pH 4 has _____ times as many hydrogen ions (H⁺) as a solution at pH 9.
- **4. WHAT IF?** How would properties of the water molecule be affected if O and H had the same electronegativity?
- 5. INTERPRET THE DATA The concentration of the appetite-regulating hormone ghrelin is about 1.3×10^{-10} M in the blood of a fasting person. How many molecules of ghrelin are in 1 L of blood?

 For selected answers, see Appendix A.

Scientific Skills Exercise

Interpreting a Scatter Plot with a Regression Line


How Does the Carbonate Ion Concentration of Seawater Affect the Calcification Rate of a Coral Reef? Scientists predict that acidification of the ocean due to higher levels of atmospheric CO_2 will lower the concentration of dissolved carbonate ions, which living corals use to build calcium carbonate reef structures. In this exercise, you will analyze data from a controlled experiment that examined the effect of carbonate ion concentration ($[CO_3^{2-}]$) on calcium carbonate deposition, a process called calcification.

How the Experiment Was Done For several years, scientists conducted research on ocean acidification using a large coral reef aquarium at Biosphere 2 in Arizona. They measured the rate of calcification by the reef organisms and examined how the calcification rate changed with differing amounts of dissolved carbonate ions in the seawater.

Data from the Experiment The black data points in the graph form a scatter plot. The red line, known as a linear regression line, is the best-fitting straight line for these points. These data are from one set of experiments, in which the pH, temperature, and calcium ion concentration of the seawater were held constant.

INTERPREDITE DATA

When presented with a graph, first determine what each axis represents. (a) In words, what is shown on the x-axis? (Include the units.)
 (b) What is on the y-axis? (c) Which variable is the independent variable—the one that was manipulated by the researchers? (d) Which is the dependent variable—the one that responded to the treatment, which was measured by the researchers? (For additional information about graphs, see the Scientific Skills Review in Appendix F.)



Data from C. Langdon et al., Effect of calcium carbonate saturation state on the calcification rate of an experimental coral reef, *Global Biogeochemical Cycles* 14:639–654 (2000).

- 2. Based on the data shown in the graph, describe in words the relationship between carbonate ion concentration and calcification rate.
- If the seawater CO₃²⁻ concentration is 270 µmol/kg, estimate the rate of calcification and how many days it would take 1 square meter of reef to accumulate 30 mmol of CaCO₃.
- 4. (a) Which step of the process in Figure 2.25 is measured in this experiment? (b) Are the results of this experiment consistent with the hypothesis that increased atmospheric [CO₂] will slow the growth of coral reefs? Why or why not?
- New Instructors: A version of this Scientific Skills Exercise can be assigned in Mastering Biology.



Chapter Review

Si Go to Mastering Biology for Assignments, the eText, the Study Area, and Dynamic Study Modules.

SUMMARY OF KEY CONCEPTS

To review key terms, go to the Vocab Self-Quiz in the Mastering Biology eText or Study Area, or go to goo.gl/ywgNry.

CONCEPT 2.1


Matter consists of chemical elements in pure form and in combinations called compounds (pp. 23-24)

- · Elements cannot be broken down chemically to other substances. A compound contains two or more different elements in a fixed ratio. Oxygen, carbon, hydrogen, and nitrogen make up approximately 96% of living matter.
- ② Compare an element and a compound.

CONCEPT 2.2

An element's properties depend on the structure of its atoms (pp. 24-27)

· An atom, the smallest unit of an element, has the following components:



- · An electrically neutral atom has equal numbers of electrons and protons; the number of protons determines the atomic number. Isotopes of an element differ from each other in neutron number and therefore mass. Unstable isotopes give off particles and energy as radioactivity.
- In an atom, electrons occupy specific electron shells; the electrons in a shell have a characteristic energy level. Electron distribution in shells determines the chemical behavior of an atom. An atom that has an incomplete outer shell, the valence shell, is reactive.

DRAW IT Draw the electron distribution diagrams for neon (10Ne) and argon (18Ar). Why are they chemically unreactive?

CONCEPT 2.3

The formation and function of molecules depend on chemical bonding between atoms (pp. 27-32)

- · Chemical bonds form when atoms interact and complete their valence shells. Covalent bonds form when pairs of electrons are shared. H₂ has a **single bond**: H—H. A **double bond** is the sharing of two pairs of electrons, as in O=O.
- Molecules consist of two or more covalently bonded atoms. The attraction of an atom for the electrons of a covalent bond is its electronegativity. Electrons of a polar covalent bond are pulled closer to the more electronegative atom, such as the oxygen in H2O.
- · An ion forms when an atom or molecule gains or loses an electron and becomes charged. An ionic bond is the attraction between two oppositely charged ions, such as Na+ and Cl-.
- · Weak interactions reinforce the shapes of large molecules and help molecules adhere to each other. A hydrogen bond is an attraction between a hydrogen atom carrying a partial positive charge (δ+) and an electronegative atom carrying a partial negative charge $(\delta -)$.

Van der Waals interactions occur between transiently positive and negative regions of molecules.

- · Molecular shape is usually the basis for the recognition of one biological molecule by another.
- 2 In terms of electron sharing between atoms, compare nonpolar covalent bonds, polar covalent bonds, and the formation of ions.

CONCEPT 2.4

Chemical reactions make and break chemical bonds (pp. 32-33)

- Chemical reactions change reactants into products while conserving matter. All chemical reactions are theoretically reversible. Chemical equilibrium is reached when the forward and reverse reaction rates are equal.
- What would happen to the concentration of products if more reactants were added to a reaction that was in chemical equilibrium? How would this addition affect the equilibrium?

CONCEPT 2.5

Hydrogen bonding gives water properties that help make life possible on Earth (pp. 33-41)

· A hydrogen bond forms when a partially negatively charged region on the oxygen of one water molecule is attracted to the partially positively charged hydrogen of a nearby water molecule. Hydrogen bonding between water molecules is the basis for water's properties.

· Hydrogen bonding keeps water molecules close to each other, giving water cohesion. Hydrogen bonding is also responsible for water's surface tension.

· Water has a high specific heat: Heat is absorbed when hydrogen bonds break and is released when hydrogen bonds form. This helps keep temperatures relatively steady, within limits that permit life. Evaporative cooling is based on water's high heat of vaporization. The evaporative loss of the most energetic water molecules cools a surface.

· Ice floats because it is less dense than liquid water. This property allows life to exist under the frozen surfaces of lakes and seas.

· Water is an unusually versatile solvent because its polar molecules are attracted to ions and polar substances that



Ice: stable hydrogen bonds




Liquid water: transient hydrogen

can form hydrogen bonds. Hydrophilic substances have an affinity for water; hydrophobic substances do not. Molarity, the number of moles of solute per liter of solution, is used as a measure of solute concentration in solutions. A mole is a certain number (6.02 \times 10²³) of molecules of a substance. The mass of a mole of a substance in grams is the same as the molecular mass in daltons.

 A water molecule can transfer an H⁺ to another water molecule to form H₃O⁺ (represented simply by H⁺) and OH⁻.

- The concentration of
 H⁺ is expressed as **pH**;
 pH = -log [H⁺]. A **buffer** consists of an acid-base pair
 that combines reversibly with
 hydrogen ions, allowing it to
 resist pH changes.
- The burning of fossil fuels increases the amount of CO₂ in the atmosphere. Some CO₂ dissolves in the oceans, causing ocean acidification, which has potentially grave conse-



quences for marine organisms that rely on calcification.

Describe how the properties of water result from the molecule's polar covalent bonds and how these properties contribute to Earth's suitability for life.

TEST YOUR UNDERSTANDING

Solution For more multiple-choice questions, go to the Practice Test in the Mastering Biology eText or Study Area, or go to goo.gl/BghzNr.

LEVELS 1-2: Remembering/Understanding

- 1. The reactivity of an atom arises from
 - (A) the average distance of the outermost electron shell from the nucleus.
 - (B) the existence of unpaired electrons in the valence shell.
 - (C) the sum of the potential energies of all the electron shells.
 - (D) the potential energy of the valence shell.
- **2.** Which of the following statements correctly describes any chemical reaction that has reached equilibrium?
 - (A) The concentrations of products and reactants are equal.
 - (B) The reaction is now irreversible.
 - (C) Both forward and reverse reactions have halted.
 - (D) The rates of the forward and reverse reactions are equal.
- **3.** Many mammals control their body temperature by sweating. Which property of water is most directly responsible for the ability of sweat to lower body temperature?
 - (A) water's change in density when it condenses
 - (B) water's ability to dissolve molecules in the air
 - (C) the release of heat by the formation of hydrogen bonds
 - (D) the absorption of heat by the breaking of hydrogen bonds
- **4.** We can be sure that a mole of table sugar and a mole of vitamin C are equal in their
 - (A) mass.

(C) number of atoms.

(B) volume.

- (D) number of molecules.
- 5. Measurements show that the pH of a particular lake is 4.0. What is the hydrogen ion concentration of the lake?
 - (A) 4.0 M
- (B) 10⁻⁴ M
- (C) $10^4 M$
- (D) 10⁻¹⁰ M

LEVELS 3-4: Applying/Analyzing

- 6. The atomic number of sulfur is 16. Sulfur combines with hydrogen by covalent bonding to form a compound, hydrogen sulfide. Based on the number of valence electrons in a sulfur atom, predict the molecular formula of the compound.
 - (A) HS
- (B) HS₂
- $(C) H_2S$
- (D) H_3S_2
- 7. What coefficients must be placed in the following blanks so that all atoms are accounted for in the products?

 $C_6H_{12}O_6 \rightarrow __C_2H_6O + __CO_2$

- (A) 1; 2
- (B) 3; 1
- (C) 1; 3
- (D) 2; 2

- 8. A slice of pizza has 500 kcal. If we could burn the pizza and use all the heat to warm a 50-L container of cold water, what would be the approximate increase in the temperature of the water?

 (Note: A liter of cold water weighs about 1 kg.)

 (A) 50°C

 (B) 5°C

 (C) 1°C

 (D) 10°C
- DRAW IT Draw the hydration shells that form around a potassium ion and a chloride ion when potassium chloride (KCl) dissolves in water. Label the positive, negative, and partial charges on the atoms.
- **10. MAKE CONNECTIONS** What do climate change (see Concept 1.1) and ocean acidification have in common?

LEVELS 5-6: Evaluating/Creating

11. SCIENTIFIC INQUIRY

Female luna moths (Actias luna) attract males by emitting chemical signals that spread through the air. A male hundreds of meters away can detect these molecules and fly toward their source. The sensory organs responsible for this behavior are the comblike antennae visible in the photograph shown here. Each filament of an antenna is equipped with thousands of receptor cells that detect



the sex attractant. (a) Based on what you learned in this chapter, propose a hypothesis to account for the ability of the male moth to detect a specific molecule in the presence of many other molecules in the air. (b) Describe predictions your hypothesis enables you to make. (c) Design an experiment to test one of these predictions.


12. FOCUS ON A THEME: EVOLUTION

The percentages of naturally occurring elements making up the human body are similar to the percentages of these elements found in other organisms. How could you account for this similarity among organisms? Explain your thinking.

13. FOCUS ON A THEME: ORGANIZATION

Several emergent properties of water contribute to the suitability of the environment for life. In a short essay (100–150 words), describe how the ability of water to function as a versatile solvent arises from the structure of water molecules.

14. SYNTHESIZE YOUR KNOWLEDGE



How do cats drink? Using high-speed video, scientists have shown that cats use an interesting technique to drink aqueous substances like water and milk. Four times a second, the cat touches the tip of its tongue to the water and draws a column of water up into its mouth (as you can see in the photo), which then shuts before gravity can pull the water back down. Describe how

the properties of water allow cats to drink in this fashion, including how water's molecular structure contributes to the process.

For selected answers, see Appendix A.

Explore Scientific Papers with Science in the Classroom MAAAS
How are some coral reefs responding to climate change?
Go to "Take the Heat" at www.scienceintheclassroom.org.
Instructors: Questions can be assigned in Mastering Biology.