

Recommended Immunization Sched. 0-18 yrs.	Birth	2mo.	4mo.	6mo.	12mo.	15 mo.	18mo.	2yr	5yr	10-12yr	16-18yr	
Hep B	X	(X	X	X)								Combined @2,4,6 mo
Dtap		(X	X	X)			X		(X)*			(Pedarix)
Polio		(X	X	X)					(X)*			Combined @ 5yr (Kinrix)
Hib (Pedvax PRP-OMP)		X	X	(*)	X							*5 yr vaccines can be given
Prevnar		X	X	X		(X)	(X)					between 4-6 yrs
Rotataq (oral)		X	X	X								
MMR					X				(X)			Combined @ 5 yrs only
Varicella					X				(X)			(Proquad)
Hep A						X		X				
HPV (Guardasil)										X*		4th dose of PCV 13 can be
Menactra										X	X	given at 15 or 18 mo.
Tdap										X	X	
Men B											X*X*	
*Notes												

Hib can be either a 3 dose (Pedvax PRP-OMP) or 4 dose series (ActHib/Pentacel PRP-T). If 4 dose would receive dose #3 at 6mo.

HPV If initiated prior to age 15. Only 2 doses are needed with dose 1 and 2 being 6-12 months apart.

If series is initiated after 15th birthday will need 3 doses 0,2mo, and 6mo.

Men B 2 doses given to those age 16-23 years. Give 2 doses atleast 1 month apart