## **Vaccines Required for Child Care**



	Hepatitis B	<b>DTaP</b> (Diphtheria, Tetanus, Pertussis)	<b>Hib</b> (Haemophilus influenzae type B)	Polio	PCV (Pneumococcal Conjugate)	MMR (Measles, mumps rubella)	Varicella (Chickenpox)
By 3 Months	2 doses	1 dose	1 dose	1 dose	1 dose		
By 5 Months	2 doses	2 doses	2 doses	2 doses	2 doses	given before 12 g	Not routinely given before 12 months of age
By 7 Months	2 doses	3 doses	2 or 3 doses (depending on vaccine)	2 doses	3 doses		
By 16 Months	2 doses	3 doses	3 or 4 doses (depending on vaccine)	2 doses	4 doses	1 dose	1 dose
By 19 Months	3 doses	4 doses	3 or 4 doses (depending on vaccine)	3 doses	4 doses	1 dose	1 dose
By 7 years or preschool/ school entry at $\geq$ 4 years*	3 doses	5 doses	Not routinely given to children age 5 years and older	4 doses	Not routinely given to children age 5 years and older	2 doses	2 doses

## \*Children attending Preschool-12th grade must meet the immunization requirements for their grade in school.

Find the Preschool-12th grade requirement chart and in the Individual Vaccine Requirements Summary immunization requirements section of the web page: www.doh.wa.gov/SCCI See the Minimum Age and Interval Table on page 2 for required minimum age and spacing information of vaccine doses.

Find information on other vaccines that are recommended, but not required, for child care/preschool attendance at: www.immunize.org/cdc/schedules.

To request this document in another format, call 1-800-525-0127.

Deaf or hard of hearing customers, please call 711 (Washington Relay) or email civil.rights@doh.wa.gov.

Vaccine	Dose #	Minimum Age*	Minimum Interval* Between Doses	Notes			
Hepatitis B (HepB)	Dose 1	Birth	4 weeks between dose 1 & 2	The final dose in the series should be given at least 24 weeks of age.			
	Dose 2	4 weeks	8 weeks between dose 2 & 3				
	Dose 3	24 weeks	16 weeks between dose 1 & 3				
Diphtheria, Tetanus, and Pertussis (DTaP)	Dose 1	6 weeks	4 weeks between dose 1 & 2	Typical vaccine schedule: 2, 4, 6 and 15-18 months of age.			
	Dose 2	10 weeks	4 weeks between dose 2 & 3	Recommended: 6 months between dose 3 and 4, but at least 4 months minimum interval			
	Dose 3	14 weeks	6 months between dose 3 & 4	acceptable.			
	Dose 4	12 months	6 months between dose 4 & 5				
	Dose 5	4 years	_				
Haemophilus influenzae	Dose 1	6 weeks	4 weeks between dose 1 & 2	If all 3 doses of PedvaxHIB given, only need 3 doses total. Dose 3 must be ≥12 months of			
type B (Hib)	Dose 2	10 weeks	4 weeks between dose 2 & 3	age.			
	Dose 3	14 weeks	8 weeks between dose 3 & 4	Only one dose required if the dose is given on or after 15 months of age. Review the Individual Vaccine Requirements Summary for minimum doses required: <a href="https://">https://</a>			
	Dose 4	12 months	_	www.doh.wa.gov/SCCI			
Pneumococcal Conjugate (PCV13)	Dose 1	6 weeks	4 weeks between dose 1 & 2	Only one dose is required if the dose is given on or after 24 months of age. Review the			
	Dose 2	10 weeks	4 weeks between dose 2 & 3	Individual Vaccine Requirements Summary for minimum doses required: <a href="https://www.doh.wa.gov/SCCI">https://www.doh.wa.gov/SCCI</a>			
	Dose 3	14 weeks	8 weeks between dose 3 & 4				
	Dose 4	12 months	_				
Polio (IPV or OPV)	Dose 1	6 weeks	4 weeks between dose 1 & 2	Three doses are acceptable if the child received dose 3 on or after their 4th birthday.			
	Dose 2	10 weeks	4 weeks between dose 2 & 3				
	Dose 3	14 weeks	6 months between dose 3 & 4				
	Dose 4	4 years	-				
Measles, Mumps, and Rubella (MMR or MMRV)	Dose 1	12 months	4 weeks between dose 1 & 2	MMRV (MMR + Varicella) may be used in place of separate MMR and varicella vaccines.			
	Dose 2	13 months	_	Must be given the same day as varicella OR at least 28 days apart, also see* footnote.			
Varicella (Chickenpox) (VAR)	Dose 1	12 months	3 months between dose 1 & 2 (12 months through 12 years). 4 weeks between dose 1 & 2 (13 years and older)	Recommended: 3 months between varicella doses, but at least 28 days minimum interval is acceptable.  Must be given the same day as MMR OR at least 28 days apart, also see* footnote.			
	Dose 2	15 months	_				

<sup>\*</sup>The 4 day grace period can be applied to all doses except between two doses of different live vaccines (such as MMR, MMRV, varicella, and Flumist).

To request this document in another format, call 1-800-525-0127.

Deaf or hard of hearing customers, please call 711 (Washington Relay) or email civil.rights@doh.wa.gov.