

Anaphylaxis Emergency Action Plan

Patient Name:		Age:
Asthma Yes (high risk for sever	re reaction) 🔲 No	
Additional health problems besides	anaphylaxis:	
Concurrent medications:		
MOUTH THROAT* SKIN GUT LUNG* HEART*	Symptoms of Anaphylaxis itching, swelling of lips and/or to itching, tightness/closure, hoarse itching, hives, redness, swelling vomiting, diarrhea, cramps shortness of breath, cough, whee weak pulse, dizziness, passing or	eness
	may be present. Severity of sympto symptoms can be life-threatening. A	
Emergency Action Steps - D	O NOT HESITATE TO GIVE EPINEPI	HRINE!
1. Inject epinephrine in thigh usin	g (check one): 🔲 Auvi-Q (0.15 mg)	Auvi-Q (0.3 mg)
	☐ EpiPen Jr (0.15 n	ng) 🗌 EpiPen (0.3 mg)
	Other (0.15 mg)	Other (0.3 mg)
Specify others:		
IMPORTANT: ASTHMA INHALERS	AND/OR ANTIHISTAMINES CAN'T B	E DEPENDED ON IN ANAPHYLAXI
2. Call 911 or rescue squad (before	calling contact)	•
3. Emergency contact #1: home	work	cell
Emergency contact #2: home	work	cell
Emergency contact #3: home	work work	cell
omments:		
		A ³
· · · · · · · · · · · · · · · · · · ·		···
octor's Signature/Date/Phone Numb	per	
arent's Signature (for Individuals ur		