MAINE ASTHMA ACTION / MANAGEMENT PLAN Name: Date of Birth: Personal Best / Predicted Peak Flow: Symptoms: Action to Take: GREEN ZONE PEAK FLOW = --☐ Continue to take your regular controller medicines every day (see blue box below). You are doing great if: ☐ Controller medicine is not needed · You aren't coughing, wheezing or having difficulty breathing Use your quick relief medication every 4-6 hours if needed for symptoms of cough, wheeze, You can sleep through the night without shortness of breath or dropping peak flows (see yellow zone) waking up with cough · You can do your usual activities ☐ Exercise pre-treatment: Take your quick relief inhaler 10-15 · Your peak flow is 80-100% of personal minutes before exercise (fill in name of inhaler and # of puffs) □ No exercise pre-treatment needed Avoid your triggers: YELLOW ZONE PEAK FLOW = Keep taking your controller medicines. START QUICK RELIEF MEDICATION: (\(\sigma\) appropriate box (es); specify dose) Your asthma is getting worse if: Make sure that your inhaler is primed first use a spacer/ chamber ☐ Xopenex MDI ___puffs ☐ Albuterol MDI puffs Every 4-6 hours as needed · You are coughing, wheezing, short of ☐ Xopenex neb ☐ Albuterol neb mg mg breath, and using quick relief medicine more than 2 extra times per week ☐ Other: · You are waking at night due to cough or wheeze more than 2 times a month · You can't do regular activities · Your peak flow is 50-80% of personal *If AT SCHOOL, give the quick relief inhaler, then CALL PARENT; may repeat medicine in best 10 minutes if not back into green zone. *If quick relief medicine is not working or you are not getting better in 24-48 hours, call your healthcare provider. RED ZONE: GET HELP NOW if: PEAK FLOW < Take a nebulizer treatment or 4 puffs of quick relief inhaler medicine now · You are very short of breath You have a hard time walking or talking → If at school, also notify parent · Skin in your neck or between ribs pulls in Call your healthcare provider now or go to the emergency department OR Call 911 Your quick relief medicine is not helping Your peak flow < 50% of personal best Other instructions: Controller Medications for Persistent Asthma: Dose Frequency Controller Medication □ 0.25mg □ 0.5mg ☐ 1.0mg times/day ☐ Budesonide Respules ☐ Pulmicort Flexhaler ☐ 90mcg ☐ 180 mcg puffs times/day ☐ 50mcg diskus ☐ 44mcg Use your regular preventive controller ☐ Fluticasone (Flovent) puffs times/day □ 110mcg □ 220mcg medication EVERY DAY as prescribed □ 10mg At bedtime ☐ Montleukast (Singulair) ☐ 4mg □ 5ma by your doctor. This will help your ☐ Asmanex 220 mcg puffs times/day asthma stay in control by decreasing the ☐ Symbicort □ 80/4.5 □160/4.5 2 puffs twice daily number of asthma flares and by □ 500/50 □ 100/50 □ 250/50 1 puff twice daily ☐ Advair diskus improving your overall lung health. □ Other: If patient is a student in school or daycare: Parent / Guardian Phone Numbers: TO BE COMPLETED BY PARENT / GUARDIAN: My child may carry and use his / her: Epi-Pen ☐ Yes ☐ No ☐ N/A Inhaled Asthma Medicine ☐ Yes ☐ No I authorize the exchange of medical information about my child's asthma between the physician's office and school nurse. DATE: PARENT / GUARDIAN SIGNATURE: TO BE COMPLETED BY PHYSICIAN / HEALTHCARE PROVIDER:

NO changes from previous plan Inhaled Medication ☐ Yes ☐ No Epi-Pen ☐ Yes ☐ No This student has the knowledge to carry and use: Please contact healthcare provider and parent if student is using quick relief medicine more than 2 times a week (i.e. in excess of pre-exercise treatment) PHONE #: FAX #: HEALTHCARE PROVIDER NAME : HEALTH CARE PROVIDER SIGNATURE: DATE: TO BE COMPLETED BY SCHOOL NURSE: Maine law now permits students to carry and use inhaled medications and epi-pen after demonstrating appropriate use to school nurse. This student demonstrates knowledge / skill to carry and use: Quick Relief Inhaler 🗆 Yes 🗀 No Epi-Pen 🗆 Yes 🗀 No SCHOOL NURSE SIGNATURE SCHOOL NAME: REVISED 11/30/07 PHONE #: EMAIL : _ FAX #: