

For Parents - Vaccines Required to Attend School in Oklahoma 2014-15 School Year

VACCINES	GRADE	PRE-SCHOOL/ PRE-K	KG-6 th	7 th – 10 th	11 th -12 th
DTP/DTaP (diphtheria, tetanus, pertussis)		4 DTaP	5 DTP/DTaP*	5 DTP/DTaP* & 1 Tdap [♦]	5 DTP/DTaP*
IPV/OPV (inactivated polio/oral polio)		3 IPV/OPV	4 IPV/OPV [◀]	4 IPV/OPV [◀]	4 IPV/OPV [◀]
MMR (measles, mumps, rubella)		1 MMR	2 MMR	2 MMR	2 MMR
Hep B (hepatitis B)		3 Hep B	3 Hep B [■]	3 Hep B [■]	3 Hep B [■]
Hep A (hepatitis A)		2 Hep A	2 Hep A	2 Hep A	2 Hep A
Varicella (chickenpox)		1 Varicella	1 Varicella	1 Varicella	1 Varicella

★	If the 4th dose of DTP/DTaP is administered on or after the child's 4th birthday, then the 5th dose of DTP/DTaP is not required.
♦	Tdap (tetanus, diphtheria, pertussis) booster
◀	If the 3rd dose of IPV/OPV is administered on or after the child's 4th birthday, then the 4th dose of IPV/OPV is not required.
■	If an adolescent reaches age 11 years and has not started the HepB vaccine series, he or she may receive a 2-dose series of Merck® Adult Hepatitis B vaccine instead of the 3-dose series of pediatric HepB. However, the series must be completed before the 16th birthday or the adolescent must receive a total of 3 doses of HepB vaccine. If you have any questions about this 2-dose series of HepB vaccine, talk to your healthcare provider. All other children (younger or older) must receive 3 doses of hepatitis B vaccine.

Vaccines Recommended for All Children but Not Required by Oklahoma School Law

A 2nd dose of varicella (chickenpox) vaccine is recommended at 4-6 years of age and any age after that if it is missed.

One dose of MCV4 (meningococcal vaccine) is recommended at age 11-12 years with a booster dose at 16 years of age.

- If an adolescent misses MCV4 at 11-12 years catch-up vaccination is recommended for ages 13 -18 years.
- If the first dose is given at age 13 through 15 years, a booster dose should be administered at age 16 through 18 years with at least 8 weeks between doses.

One dose of MCV4 (meningococcal vaccine) is recommended for first-year college students up through age 21 years who are living in residence halls if they have not received a dose on or after their 16th birthday. Oklahoma law requires a dose of MCV4 for first time college enrollees who reside in on-campus student housing.

Three doses of HPV (human papillomavirus) vaccine are recommended starting at 11-12 years of age.

- If an adolescent misses HPV at 11-12 years catch-up vaccination is recommended for females 13-26 years and for males 13 -21 years.

Vaccine-Preventable Diseases and the Vaccines that Prevent Them

Diphtheria (Can be prevented by DTaP & Tdap vaccines)

Diphtheria is a very contagious bacterial disease that affects the respiratory system, including the lungs. Diphtheria bacteria can be passed from person to person by direct contact with droplets when an infected person coughs or sneezes. When people are infected, the diphtheria bacteria produce a toxin (poison) in the body that can cause weakness, sore throat, low-grade fever, and swollen glands in the neck. Effects from this toxin can also lead to swelling of the heart muscle and, in some cases, heart failure. In severe cases, the illness can cause coma, paralysis, and even death.

Hepatitis A (Can be prevented by HepA vaccine)

Hepatitis A is an infection of the liver caused by hepatitis A virus. The virus is usually spread person-to-person through the fecal-oral route. In other words, the virus is taken in by mouth from contact with objects, food, or drinks contaminated by the feces (stool) of an infected person. Symptoms include fever, tiredness, loss of appetite, nausea, abdominal discomfort, dark urine, and jaundice (yellowing of the skin and eyes). An infected person may have no symptoms, may have mild illness for a week or two, or may have severe illness for several months that requires hospitalization. In the U.S. about 100 people a year die from hepatitis A.

Hepatitis B (Can be prevented by HepB vaccine)

Hepatitis B is an infection of the liver caused by hepatitis B virus. The virus spreads through exposure to blood or other body fluids, for example, from sharing personal items, such as razors or during sex. Hepatitis B causes a flu-like illness with loss of appetite, nausea, vomiting, rashes, joint pain, and jaundice. The virus stays in the liver of some people for the rest of their lives and can result in severe liver diseases, including fatal cancer.

Human Papillomavirus (Can be prevented by HPV vaccine)

Human Papillomavirus also known as HPV, is a very common virus that is spread by skin-to-skin contact during any type of sexual activity with another person. About 79 million Americans, most in their late teens and early 20s, are infected with HPV. HPV is so common that nearly all sexually active men and women get it at some point in their lives. It is a major cause of cervical cancer in women and genital warts in women and men. Every year in the United States about 4,000 women die from cervical cancer caused by HPV and about 8,000 cancers caused by HPV occur each year in men.

Measles (Can be prevented by MMR vaccine)

Measles is one of the most contagious viral diseases. Measles virus is spread by direct contact with the airborne respiratory droplets of an infected person. Measles is so contagious that just being in the same room after a person who has measles has already left can result in infection. Symptoms usually include a rash, fever, cough, and red, watery eyes. Fever and rash can last for up to a week, and the coughing lasts about 10 days. Measles can lead to pneumonia, seizures, brain damage, and death.

Meningococcal Disease (Can be prevented by MCV vaccine)

Meningococcal disease is caused by bacteria and is a leading cause of bacterial meningitis (infection around the brain and spinal cord) in children. The bacteria are spread by droplets from the nose and throat through coughing, sneezing or kissing. Symptoms include nausea, vomiting, sensitivity to light, confusion and sleepiness. Meningococcal disease also causes blood infections. About one of every ten people who get the disease dies from it. Survivors of meningococcal disease may lose their arms or legs, become deaf, have problems with their nervous systems, become developmentally disabled, or suffer seizures or strokes.

Mumps (Can be prevented by MMR vaccine)

Mumps is an infectious disease caused by the mumps virus, which is spread in the air by a cough or sneeze from an infected person. A child can also get infected with mumps by coming in contact with a contaminated object, like a toy. The mumps virus causes fever, headaches, painful swelling of the salivary glands under the jaw, muscle aches, tiredness, and loss of appetite. Severe complications for children who get mumps are uncommon, but can include meningitis (infection of the covering of the brain and spinal cord), encephalitis (inflammation of the brain), permanent hearing loss, or swelling of the testes, which rarely can lead to sterility in men.

Pertussis (Whooping Cough) (Can be prevented by DTaP & Tdap vaccines)

Pertussis is caused by bacteria spread through direct contact with respiratory droplets when an infected person coughs or sneezes. In the beginning, symptoms of pertussis are similar to the common cold, including runny nose, sneezing, and cough. After 1-2 weeks, pertussis can cause spells of violent coughing and choking, making it hard to breathe, drink, or eat. This cough can last for weeks. Pertussis is most serious for babies, who can get pneumonia, have seizures, become brain damaged, or even die. About two-thirds of

children under 1 year of age who get pertussis must be hospitalized.

Polio (Can be prevented by IPV vaccine)

Polio is caused by a virus that lives in an infected person's throat and intestines. It spreads through contact with the feces (stool) of an infected person and through droplets from a sneeze or cough. Symptoms typically include sudden fever, sore throat, headache, muscle weakness, and pain. In about 1% of cases, polio can cause paralysis. Among those who are paralyzed, up to 5% of children may die because they become unable to breathe.

Rubella (German Measles) (Can be prevented by MMR vaccine)

Rubella is caused by a virus that is spread through coughing and sneezing. In children rubella usually causes a mild illness with fever, swollen glands, and a rash that lasts about 3 days. Rubella rarely causes serious illness or complications in children, but can be very serious to a baby in the womb. If a pregnant woman is infected, the result to the baby can be devastating, including miscarriage, serious heart defects, mental retardation, and loss of hearing and eye sight.

Tetanus (Lockjaw) (Can be prevented by Tdap vaccine)

Tetanus is caused by bacteria found in soil. The bacteria enter the body through a wound, such as a deep cut. When people are infected, the bacteria produce a toxin (poison) in the body that causes serious, painful spasms and stiffness of all muscles in the body. This can lead to "locking" of the jaw so a person cannot open his or her mouth, swallow, or breathe. Complete recovery from tetanus can take months. Three of ten people who get tetanus die from the disease.

Varicella (Chickenpox) (Can be prevented by varicella vaccine)

Chickenpox is caused by the varicella zoster virus. Chickenpox is very contagious and spreads very easily from infected people. The virus can spread from either a cough or a sneeze. It can also spread from the blisters on the skin, either by touching them or by breathing in the viral particles. Typical symptoms of chickenpox include an itchy rash with blisters, tiredness, headache and fever. Chickenpox is usually mild, but it can lead to severe skin infections, pneumonia, encephalitis (brain swelling), and even death.